


Research team confirms icy conditions existed in South Pole region during Late Cretaceous period

September 7 2022, by Sonia Fernandez



Bedrock map of Northern and Southern Victoria Land, Antarctica, showing the distribution of outcrop and location of the Butcher Ridge Igneous Complex (BRIC) and other localities referred to in the text with demonstrated mid to late-Cretaceous alteration ages. Basemap constructed using Quantarctica v3.2 from the Norwegian Polar Institute (Matsuoka, K. et al. Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic

islands. Environmental Modeling and Software 140, 105015 (2021)) with Rock outcrop data from SCAR Antarctic Digital Database (ADD) Version 7.0. Credit: Demian A. Nelson et al, *Nature Communications* (2022). DOI: 10.1038/s41467-022-32736-9

Let's pretend it's the Late Cretaceous, roughly 66 to 100 million years ago. We've got dinosaurs roaming the land and odd-looking early species of birds, although the shark as we know it is already swimming in the prehistoric oceans—which cover 82% of Earth. Redwood trees and other conifers are making their debut, as are roses and flowering plants, and with them come bees, termites and ants. Most of all, it's warm, volcanically active and humid all over the place with nary an ice sheet in sight.

Except, according to a group of scientists from UC Santa Barbara, University of Oregon and University of Manitoba, icy conditions did exist in the region of the South Pole.

"And it wasn't just a single-valley glacier," said UCSB geologist John Cottle, "it was probably multiple glaciers or a large [ice sheet](#)." Contrary to our widely held picture of the Late Cretaceous as "hot everywhere," he said, there's evidence that polar ice existed during that period, even at the height of global greenhouse conditions. The geologists' study is published in the journal *Nature Communications*.

A prehistoric puzzle

Fast-forward to today. Let's pretend we're in Antarctica. It's chilly, it's barren, and we're standing near a large grouping of exposed glassy rock along the Transantarctic Mountains, adjacent to the Ross Ice Shelf, called the Butcher Ridge Igneous Complex (BRIC).

"I actually heard about these rocks when I was a grad student 20 or so years ago, and they're just really weird," Cottle said. Remote, even by today's Antarctic exploration standards, the BRIC is unusual because the rocks' composition and formation are uncharacteristic of nearby rock formations, with—among other things—large amounts of glass and layered alteration that indicates significant physical, chemical or environmental events that changed their mineral composition.

Cottle got the chance to finally sample the BRIC on a recent expedition, and in the process of analyzing how it was formed, he and his team encountered an "unusually large amount of water."

"So you have a really hot rock that interacts with water, and as it cools, incorporates it into the glass," he said. "If you look at the composition, then you can tell something about where that water came from. It can exist as hydroxyl, which tells you that it probably came from the [magma](#), or it could be molecular, which means it is probably external."

What they were expecting to see was that the alteration in the rock was caused by the water already in the magma as it cooled. What they found instead was a record of a climate process that was thought not to have existed at the time.

In their spectroscopic analysis of the samples, the researchers determined that while some of the water indeed originated with magma as it plumed upward from Earth's interior, as the molten rock cooled into glass just beneath the Earth's surface, it also incorporated groundwater.

"We determined that most of the water in these rocks is externally derived," Cottle said. "We then measured the oxygen and hydrogen isotopic composition of the water and it matches very well to the composition of Antarctic snow and ice."

To lock in their result, Cottle and team also conducted argon-argon geochronology to date the rock and its alteration.

"The problem is, these rocks are Jurassic, so about 183 million years old," he said. "So when you measure the alteration, what you don't know is when that happened." They were able to recover the age of the rock (Jurassic), but also found a younger age (Cretaceous). "So when these rocks cooled and were altered," he continued, "it also reset the argon isotope as well, and you can match the age of the alteration to the composition of the alteration."

There are other, similar volcanic rocks roughly 700 km north of the BRIC that also have a Cretaceous alteration age, indicating that polar glaciation might have been regionally extensive in Antarctica during that time. "What we'd like to do is go to other places in Antarctica and see if we can determine the scale of the glaciation, if we recover the same results that we've already found," he said.

Finding evidence of large ice sheets dating back to the Cretaceous might not alter our general picture of a hot and humid Earth at that time, Cottle said, "but we would have to think about the Cretaceous and Antarctica quite differently than we do now."

Research in this study was also conducted by Demian A. Nelson (lead author) of UCSB, Ilya N. Bindeman at University of Oregon and Alfredo Camacho at University of Manitoba.

More information: Demian A. Nelson et al, Ultra-depleted hydrogen isotopes in hydrated glass record Late Cretaceous glaciation in Antarctica, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-32736-9](https://doi.org/10.1038/s41467-022-32736-9)

Provided by University of California - Santa Barbara

Citation: Research team confirms icy conditions existed in South Pole region during Late Cretaceous period (2022, September 7) retrieved 28 June 2024 from <https://phys.org/news/2022-09-team-icy-conditions-south-pole.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.