


World's lakes experience excess warming

September 12 2022



Excess lake warming in the ice-off or ice-on month. a Trends in monthly lake surface water temperature (LSWT) from 1979 to 2020 across three lakes in Europe, Asia and North America, respectively. The location of the three lakes is shown as black dots in b, c. The dashed lines and shaded areas represent the mean and range of ice-off date or ice-on date from 1979 to 2020. b, c Spatial pattern of mean ice-off month (b), ice-on month (c). d Mean trend in monthly LSWT from 1979 to 2020 across lakes with the same ice-off month from March to July. e, f Spatial pattern of ratios of the trend in LSWT in the ice-off month ($\text{LSWT}_{\text{IOFF}}$) or LSWT in the ice-on month (LSWT_{ION}) to the trend in LSWT during the open-water period (RIOFF (e) and RION (f)). The inset in b, c, e, f shows the probability density function (PDF) from Europe (orange), North

America (pink) and Asia (blue), respectively. Credit: *Nature Communications* (2022). DOI: 10.1038/s41467-022-32830-y

More than half the world's 117 million lakes experience ice cover for part of the year. But with the rise in global temperatures, the timing of seasonal ice formation and loss are changing at rapid rates. Since 1979, the average timing of lake ice break-up across the Northern Hemisphere has advanced by 8 days. This 8-day advancement has resulted in excess lake warming with numerous implications for lake ecosystems.

A new paper published in *Nature Communications* by Dr. Iestyn Woolway of Bangor University and colleagues at Peking University has concentrated on Northern Hemisphere lakes, using [satellite images](#) to study nearly a thousand lakes and modeling over 100,000 to calculate the effects of the increased ice-free days on the [water temperature](#).

They also suggest how changing temperature profiles within lakes might affect the processes within the lakes as well as the plants and animals living within the lake environments.

"Projecting future lake warming due to changes in ice cover is important for understanding how these changes could affect ecological processes within lakes, and how that will affect all the ways we, and other animals and plants use lakes in the future," explained Dr. Iestyn Woolway of the University's School of Ocean Sciences.

"Higher lake surface-temperatures due to shorter periods of ice cover can impact processes within the lakes, leading to effects such as increasing or earlier plankton blooms, and increase in cyanobacteria and an increase in layering of different water temperatures. Water stratification could lead to lake deoxygenation [low oxygen

concentrations] at depth, and loss of habitat for oxygen-sensitive fish," he added.

Iestyn and his fellow researchers estimate that lakes will be ice-free around a month earlier (15–45 days in the Northern Hemisphere), which may lead to an increase of between 2 and 6 degrees additional surface temperature (during the ice-off month, the time of year in which dormant species begin to thrive once again) this century. They have also estimated that each newly ice-free day adds an average increase in surface lake-temperature between 1.1–0.9°C.

More information: Xinyu Li et al, Earlier ice loss accelerates lake warming in the Northern Hemisphere, *Nature Communications* (2022). DOI: [10.1038/s41467-022-32830-y](https://doi.org/10.1038/s41467-022-32830-y)

Provided by Bangor University

Citation: World's lakes experience excess warming (2022, September 12) retrieved 26 April 2024 from <https://phys.org/news/2022-09-world-lakes-excess.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--