

Young, cold-blooded animals are suffering the most as Earth heats up, research finds

September 20 2022, by Patrice Pottier



Credit: AI-generated image (disclaimer)

Climate change is <u>making heat waves</u> worse. Many people have already noticed the difference—and so too have other animals.

Sadly, <u>research</u> by myself and colleagues has found <u>young animals</u>, in particular, are struggling to keep up with rising temperatures, likely



making them more vulnerable to <u>climate change</u> than adults of their species.

The study focused on "ectotherms," or cold-blooded <u>animals</u>, which comprise <u>more than 99%</u> of animals on Earth. They include fish, reptiles, amphibians and insects. The body <u>temperature</u> of these animals reflects outside temperatures—so they can get <u>dangerously hot</u> during <u>heat waves</u>.

In a warming world, a species' ability to adapt or acclimatize to temperatures is crucial. Our study found that young ectotherms, in particular, can struggle to handle more heat as their habitat warms up. That may have dramatic consequences for biodiversity as climate change worsens.

Our findings are yet more evidence of the need to urgently reduce greenhouse gas emissions to prevent catastrophic global heating. Humans must also provide and retain cool spaces to help animals navigate a warmer future.

Tolerating heat in a changing climate

The body temperature of ectotherms is extremely variable. As they move through their habitat, their body temperature varies according to the outside conditions.





Credit: F1lter 88 from Pexels

However, there's only so much heat these animals can tolerate. Heat tolerance is defined as the maximum body temperature ectotherms can handle before they <u>lose functions</u> such as the ability to walk or swim. During heat waves, their body temperature gets so high they can <u>die</u>.

Species, <u>including ectotherms</u>, can adapt to challenges in their environment over time by evolving across generations. But the rate at which <u>global temperatures</u> are rising means in many cases, this adaptation is not happening fast enough. That's why we need to understand how animals acclimatize to rising temperatures within a single lifetime.

Unfortunately, some young animals have little to no ability to move and seek cooler temperatures. For example, baby lizards inside eggs cannot



move elsewhere. And owing to their <u>small size</u>, juvenile ectotherms cannot move great distances.

This suggests young animals may be particularly vulnerable during intense heat waves. But we know <u>very little</u> about how young animals acclimatize to high temperatures. Our research sought to find out more.

Young animals at risk

<u>Our study</u> drew on 60 years of research into 138 ectotherm species from around the world.

Overall, we found the heat tolerance of embryos and juvenile ectotherms increased very little in response to rising temperatures. For each degree of warming, the <u>heat tolerance</u> of young ectotherms only increased by an average 0.13°C.





Credit: AI-generated image (disclaimer)

The physiology of heat acclimatization in animals is very complex and poorly understood. It appears linked to a number of factors such as <u>metabolic activity</u> and <u>proteins</u> produced by cells in response to stress.

Our research showed young land-based animals were worse at acclimatizing to heat than aquatic animals. This may be because moving to a cooler temperature on land is easier than in an aquatic environment, so land-based animals may not have developed the same ability to acclimatize to heat.

Heat tolerance can vary within a species. It can <u>depend on</u> what temperatures an animal has experienced during its lifetime and, as such, the extent to which it has acclimatized. But surprisingly, our research found past exposure to high temperatures does not necessarily help a young animal withstand future high temperatures.

Take, for example, Lesueur's velvet gecko which is found mostly along Australia's east coast. Research shows juveniles from eggs incubated in cooler nests (23.2°C) tolerated temperatures up to 40.2°C. In contrast, juveniles from warmer nests (27°C) only tolerated temperatures up to 38.7° C.

Those patterns can persist through adulthood. For example, adult male <u>mosquito fish</u> from eggs incubated to 32°C were less tolerant to heat than adult males that experienced 26°C during incubation.

These results show embryos are especially vulnerable to extreme heat. Instead of getting better at handling <u>heat</u>, warmer eggs tend to produce



juveniles and adults less capable of withstanding a warmer future.

Overall, our findings suggest young cold-blooded animals are already struggling to cope with rising temperatures—and conditions during early life can have lifelong consequences.



Credit: F1lter 88 from Pexels

What's next?

To date, most studies on the impacts of climate change have focused on adults. Our <u>research</u> suggests animals may be harmed by heatwaves long before they reach adulthood—perhaps even before they're born.

Alarmingly, this means we may have underestimated the damage climate



change will cause to biodiversity.

Clearly, it's vitally important to limit global <u>greenhouse gas emissions</u> to the extent required by the <u>Paris Agreement</u>.

But we can also act to protect species at a finer scale—by conserving habitats that allow animals to find shade and shelter during heatwaves. Such habitats include trees, shrubs, burrows, ponds, caves, logs and rocks. These places must be created, restored and preserved to help animals prosper in a warming world.

More information: Patrice Pottier et al, Developmental plasticity in thermal tolerance: Ontogenetic variation, persistence, and future directions, *Ecology Letters* (2022). <u>DOI: 10.1111/ele.14083</u>

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Young, cold-blooded animals are suffering the most as Earth heats up, research finds (2022, September 20) retrieved 2 May 2024 from <u>https://phys.org/news/2022-09-young-cold-blooded-animals-earth.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.