


Zhurong rover finds evidence of possible flooding on Mars billions of years ago

September 27 2022, by Bob Yirka



Region around the Zhurong rover landing site. a, Topographic map showing the landing site of Zhurong (red star), as well as the landing sites of the Phoenix, InSight, Curiosity, Perseverance and Viking-2 landers/rovers (orange squares).

The purple solid and dashed lines show the locations of Martian palaeoshorelines of ref. 11, roughly delineating the depositional contact of the VBF in the northern plains. b, Simplified geological map near the Zhurong landing site with data from ref. 18. Scale bar, 200 km. c, Geomorphic map of the Zhurong landing area with data from ref. 22. Scale bar, 15 km. d, Traverse of the Zhurong rover from 25 May (Sol 11) to 6 September (Sol 113) 2021 on the basemap of a Tianwen-1 High Resolution Imaging Camera image (Sol 19, 2 June 2021). The red star marks the landing site (25.066° N, 109.925° E) and the red line shows the track of the rover. Scale bar, 100 m. Relative distances to the landing site are marked alongside the track. Credit: *Nature* (2022). DOI: 10.1038/s41586-022-05147-5

A team of researchers at the Chinese Academy of Sciences, working with a colleague from Peking University, has found evidence of a layered subsurface in the Utopian Basin on Mars. In their paper published in the journal *Nature*, the group describes studying radar data from the Zhurong rover and what it revealed about the ground beneath the Utopia Planitia basin on Mars.

China's National Space Administration landed the Zhurong rover on the red planet in May of 2021. Since that time, it has been rolling across the Utopia Planitia basin—an [impact crater](#)—studying the terrain around it and the material that lies beneath. To date, the rover has traveled approximately 1,171 meters. The rover is equipped with ground-penetrating [radar](#), which has been fired continuously at the ground as the rover moves, allowing the researchers to create an underground map for depths from 3 to 10 meters. The [rover](#) also has a device that can send low-frequency radio waves into the ground at depths up to 100 meters, though its resolution is much lower than the radar.

The researchers found that there are at least two layers of material beneath the basin, neither of which are believed to be water. One of the

layers was approximately 10 to 30 meters down and the other from 30 to 80 meters down.

Neither the radar nor the [radio waves](#) are capable of discerning the difference between rock and ice or lava. Thus, the researchers have resorted to theories to explain the layers beneath the ground. They suggest it is likely that the deeper, older layer came about as smaller rocks settled onto bigger rocks during a flood approximately 3 billion years ago.

The formation they found would require the flood to have been a fast one with enough energy to carry large rocks. The researchers suggest the second layer could have come about in similar fashion—by a flood approximately 1.6 billion years ago. They note that prior research has suggested there was a lot of glacial activity during that time period. They also note that they did not find any evidence of volcanic activity that could have resulted in the creation of either layer.

More information: Chao Li et al, Layered subsurface in Utopia Basin of Mars revealed by Zhurong rover radar, *Nature* (2022). [DOI: 10.1038/s41586-022-05147-5](https://doi.org/10.1038/s41586-022-05147-5)

© 2022 Science X Network

Citation: Zhurong rover finds evidence of possible flooding on Mars billions of years ago (2022, September 27) retrieved 13 March 2024 from <https://phys.org/news/2022-09-zhurong-rover-evidence-mars-billions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.