


Tracing anthropogenically emitted carbon dioxide into the ocean

October 17 2022, by Aaron Sidder



The net atmosphere-to-ocean carbon flow rates for total carbon (and hence for ΔC), anthropogenically emitted carbon C_A , and for natural carbon (and hence for ΔC_N). Solid lines are for the case of variable air-sea exchange, while dashed

lines are for constant air-sea exchange coefficients (constant solubility and piston velocities). The light shading around lines visible for recent years indicates the half-range spread across the three versions of the circulations used. Credit: *Global Biogeochemical Cycles* (2022). DOI: 10.1029/2022GB007405

Through fossil fuel combustion, deforestation, and other industrial and agricultural activities, humans have raised global atmospheric carbon dioxide (CO₂) levels to more than 415 parts per million. That concentration represents a 135-parts-per-million increase since preindustrial times in the late 18th century. It is often assumed that this CO₂ increase consists entirely of anthropogenically emitted CO₂, but a new study challenges that assumption.

To investigate the standard narrative about the fate of anthropogenically emitted carbon, Holzer and DeVries labeled CO₂ as it was emitted and tracked it with a data-assimilated ocean circulation model. This method allowed them to partition the net changes in atmospheric and oceanic carbon inventories as either anthropogenically emitted or natural carbon. The model followed the journey of emitted carbon from 1780 to 2020 using what is called a linear labeling tracer. Researchers have used this labeling technique in other applications but never to track anthropogenic carbon.

The modeling revealed that only 45% of the increase in [atmospheric carbon](#) over the past few centuries originated as emitted carbon and that outgassed natural carbon from the ocean accounts for the other 55%. The researchers also found that the ocean has accumulated nearly twice as much emitted carbon as previously assumed.

The mechanism behind the outgassing is seawater carbonate chemistry. As dissolved [inorganic carbon](#) builds up in the ocean, the seas

increasingly leak into the atmosphere CO₂ that was dissolved in the ocean before the Industrial Revolution. By 2020, the modeling showed, the ocean had lost one preindustrial CO₂ molecule for every 2.2 emitted CO₂ molecules it absorbed.

Put another way, since 1780, the oceans have outgassed approximately 160 petagrams of natural carbon while absorbing 350 petagrams of emitted carbon, for a net increase of about 190 petagrams of carbon stored in the [ocean](#).

The researchers point out that their results do not challenge estimates of anthropogenically driven changes in global carbon inventories. Rather, they quantify where emitted [carbon](#) ends up, and they illustrate that linear labeling tracers are an effective and powerful tool for quantifying the effects of nonlinear processes in the climate system.

The research was published in *Global Biogeochemical Cycles*.

More information: Mark Holzer et al, Source-Labeled Anthropogenic Carbon Reveals a Large Shift of Preindustrial Carbon From the Ocean to the Atmosphere, *Global Biogeochemical Cycles* (2022). [DOI: 10.1029/2022GB007405](#)

This story is republished courtesy of Eos, hosted by the American Geophysical Union. Read the original story [here](#).

Provided by Eos

Citation: Tracing anthropogenically emitted carbon dioxide into the ocean (2022, October 17) retrieved 3 May 2024 from <https://phys.org/news/2022-10-anthropogenically-emitted-carbon-dioxide-ocean.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.