

Breakthrough in optical information transmission: A one-way street for optical vortices

October 20 2022



Visualisation of topology-selective Brillouin scattering effect in chiral photonic crystal fibre. Linear arrows are propagation direction and circular arrows are circular polarisation states or vortex phases. Credit: *Science Advances* (2022). DOI: 10.1126/sciadv.abq6064

Scientists at the Max Planck Institute for the Science of Light have managed for the first time to create a unidirectional device that significantly increases the quality of a special class of transmitted signals in optical communications: optical vortices.

By transmitting selective optical vortex modes exclusively



unidirectionally, the developed device largely reduces detrimental backscattering to a minimum. The scientists emphasize the great practical utility of their discovery in many <u>optical systems</u>, with applications ranging from mode division multiplexed communications and optical tweezers to vortex lasers and quantum manipulation systems.

Optical communication can be improved by increasing the amount of optical information transmitted. This can be achieved by using multiplexed channels such as using many optical wavelengths, different polarization states or multiple time slots. In the last decade, optical spatial modes, which are the eigenfields in the waveguides, are widely exploited to further improve the communication capacity due to the little crosstalk between orthogonal spatial modes.

In classical communication as well as in quantum communication, the use of vortex modes in multiplexing methods has proven to be advantageous. This special mode set possesses a helical optical phase distribution and allows an additional degree of freedom for multiplexing optical signals. Devices like vortex generators, lasers and signal amplifiers were demonstrated and are in great demand.

A limiting effect on the applicability is that there has not yet been a device that permits transmission of certain vortex modes in one direction but not the opposite one. However, just this kind of device—a so-called optical vortex isolator—is of crucial importance for the improvement of signal quality and purity. The particular difficulty in developing such a device is a fundamental principle of optics: reciprocity. It requires a symmetrical response of a transmission channel when the source and observation points are interchanged.

Researchers succeed in building an optical vortex insulator



Now, a team at the Max Planck Institute for the Science of Light led by Xinglin Zeng, Philip Russell and Birgit Stiller, achieved a breakthrough that makes this possible: They used <u>sound waves</u> that propagate only in one direction to break the light transmission reciprocity for chosen vortex modes.

The effect of so-called topology-selective Brillouin-Mandelstam scattering in chiral photonic crystal fiber allows for a unidirectional interaction of vortex-carrying <u>light waves</u> with traveling sound waves. A specific optical vortex can be strongly suppressed or amplified with a well-designed control light. The experimental results published in *Science Advances* show a significant vortex isolation rate, preventing random backscattering and signal degradation in the system.

"This is the first nonreciprocal system for vortex modes, which opens up a new perspective in nonreciprocal optics—the same physical effect can happen not only on the fundamental modes but also on higher-order modes," says Xinglin Zeng, the first author of this paper.

"The light-driven optical vortex isolator will have great impact on the applications such as <u>optical communications</u>, <u>quantum information</u> <u>processing</u>, <u>optical tweezers</u>, and fiber lasers. I find the possibility of selective manipulation of <u>vortex</u> modes solely by light and sound waves a very fascinating concept," says Birgit Stiller, the leader of the Quantum Optoacoustics Research Group.

More information: Xinglin Zeng et al, Nonreciprocal vortex isolator via topology-selective stimulated Brillouin scattering, *Science Advances* (2022). DOI: 10.1126/sciadv.abq6064

Provided by Max Planck Institute for the Science of Light



Citation: Breakthrough in optical information transmission: A one-way street for optical vortices (2022, October 20) retrieved 3 May 2024 from <u>https://phys.org/news/2022-10-breakthrough-optical-transmission-one-way-street.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.