


# Ecosystem-based fisheries management restores western Baltic fish stocks

October 14 2022



Western Baltic Sea (WBS) position in the context of the northeast Atlantic shelf region including the entire Baltic Sea region (A). The WBS ecosystem (B) is bounded by ICES subdivisions 22 and 24 (red line). Credit: *Frontiers in Marine Science* (2022). DOI: 10.3389/fmars.2022.879998

Decades of overfishing, together with nutrient pollution, rapid increase in hypoxia, ocean warming and acidification have put fish and harbor porpoise (*Phocoena phocoena*) in the western Baltic Sea at risk of collapse.

But the commercially relevant stocks of cod (*Gadus morhua*), herring (*Clupea harengus*) and sprat (*Sprattus sprattus*) can be restored and prospects for marine mammals be improved, according to a team of marine scientists from GEOMAR Helmholtz Center for Ocean Research Kiel (Germany), the German Federal Agency for Nature Conservation (Bundesamt für Naturschutz, BfN, Germany) and the Institute of Biosciences and Bioresources at the National Research Council (CNR) of Italy.

Using model simulations, the researchers tested five scenarios from no fishing to ecosystem-based fisheries management. This approach accounts for the roles of species within their ecosystem and adjusts catches accordingly to keep [fish stocks](#) in a healthy, productive and resilient condition.

A study now published in *Frontiers in Marine Science* concludes that ecosystem-based fisheries management would allow the endangered harbor porpoise population to recover and increase catches of herring and cod significantly within a decade. The [food web](#) would become less susceptible to eutrophication and climate change and, in addition, more able to support carbon sequestration than in a business-as-usual scenario that assumes today's fisheries practices continue.

The study benefits from years of data collection at GEOMAR. Building on a first prototype and a huge amount of data, the researchers now developed the first model for the western Baltic Sea that includes top predators such as harbor porpoise and seals, various fish species and other marine animals, plankton, algae and seaweeds, as well as their interactions under different scenarios. "Looking at the big picture of the food web helps to identify management options that sustain important food resources and dependant businesses," says Dr. Marco Scotti, marine ecologist at GEOMAR and CNR, lead author of the recent publication.

Ecosystem-based fisheries management would imply to stop catching juvenile cod, reduce catches of herring and sprat to half of the maximum sustainable yield—the highest possible harvest per year that can be sustained over time—and catches of adult cod and flatfish to 80 percent of the maximum sustainable yield. This approach was compared to a business-as-usual scenario characterized by average fishing mortalities for all exploited stocks during the years 2015 to 2019.

In the business-as-usual scenario, cod stocks decline slightly below the numbers of 2019 and herring stocks to almost half their 2019 sizes by 2050. Sprat and flatfish increase to some degree, suggesting a substantial regime shift. Ecosystem-based fisheries management, in contrast, would lead to an increase of almost 70 percent in cod and 50 percent in herring catches by 2050, compared to the period 2015 to 2019. Catches in flatfish would increase by almost 20 percent, but with largely reduced fishing effort and costs. Potential for [carbon sequestration](#) would be more than three times greater under ecosystem-based fisheries management compared to business-as-usual.

"The Common Fisheries Policy of Europe demands an end to overfishing by 2020 and rebuilding of healthy and resilient ecosystems thereafter," says Dr. Rainer Froese, fisheries biologist at GEOMAR and co-author of the study.

"Past and present overfishing—not climate change—was the main cause of the recent collapse of herring, cod, and profitable western Baltic fisheries in general. Continued business-as-usual would in addition push the highly endangered harbor porpoise to the brink of extinction. In contrast, ecosystem-based management would rebuild healthy stocks and fisheries and even help us to fight [climate change](#). Saving the western Baltic requires to stop fishing of cod and herring for a few years, until these stocks have recovered. During this time, fishers need to be compensated for their losses. Fishing for plaice and other flatfish can

meanwhile continue."

**More information:** Marco Scotti et al, Ecosystem-based fisheries management increases catch and carbon sequestration through recovery of exploited stocks: The western Baltic Sea case study, *Frontiers in Marine Science* (2022). [DOI: 10.3389/fmars.2022.879998](https://doi.org/10.3389/fmars.2022.879998)

Provided by Helmholtz Association of German Research Centres

Citation: Ecosystem-based fisheries management restores western Baltic fish stocks (2022, October 14) retrieved 4 June 2024 from <https://phys.org/news/2022-10-ecosystem-based-fisheries-western-baltic-fish.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.