



Researchers provide new insights into photosynthesis

October 14 2022


a



b



c



Overall structure of the PSI dimer. a, View of individual proteins from stroma. b, Arrangement of the pigments in the outline of the map: chlorophylls green (Mg yellow), luteins blue, β -carotenes red, violaxanthin purple and neoxanthin pink. c, Overall view along the membrane. Credit: *Nature Plants* (2022). DOI: 10.1038/s41477-022-01253-4

Photosynthesis is the most important basis of life on Earth. In it, plants and single-cell algae use the energy of sunlight and convert this energy into sugar and biomass. In this process, oxygen is released. Plant biotechnologists and structural biologists from the Universities of Münster (Germany) and Stockholm (Sweden) have clarified the structure of a new protein complex which catalyzes energy conversion processes in photosynthesis. This protein complex is the photosystem I, which is known as a single protein complex (monomer) in plants.

The team of researchers headed by Prof. Michael Hippler from the University of Münster and Prof. Alexey Amunts from the University of Stockholm have now shown, for the first time, that two [photosystem I](#) monomers in [plants](#) can join together as a dimer, and they describe the molecular structure of this new kind of molecular machine. The results, which have been published in the *Nature Plants* journal, provide molecular insights into the photosynthesis process to a hitherto unparalleled degree of precision. They could help to utilize more efficiently in future the reductive force (i.e., the preparedness to give up [electrons](#)) of photosystem I, for example to produce hydrogen as a source of energy.

The background: There are two photosynthesis complexes, called photosystems I and II, which work at their best in the case of light with

different wavelengths. The uptake of light energy into photosystems I and II enables electrons to be transported within the molecular photosynthetic machine, thus driving the conversion of light energy into chemical energy. In the process, electrons from photosystem I are transmitted to the protein ferredoxin.

In [green algae](#), ferredoxin can transmit electrons arising during photosynthesis to an enzyme called hydrogenase, which then produces molecular hydrogen. This [molecular hydrogen](#) is thus produced by the input of light energy, which means it is produced renewably and might be able to act as a future source of energy. The researchers asked themselves the question: How does the production of photosynthetic hydrogen relate to the structural dynamics of the monomer and dimer photosystem I?

The results in detail

The photosystem I homodimer from the green alga *Chlamydomonas reinhardtii* consists of 40 protein subunits with 118 transmembrane helices providing a structure for 568 photosynthesis pigments. Using [cryogenic electron microscopy](#), the researchers showed that the absence of subunits with the designation PsaH and Lhca2 leads to a head-to-head orientation of monomer photosystem I (PSI) and its associated light-harvesting proteins (LHCI). The light-harvesting protein Lhca9 is the key element providing for this dimerization.

In the study, the researchers define the most precisely available PSI-LHCI model to a resolution of 2.3 Ångström (one Ångström corresponds to one ten-millionth of a millimeter), including the flexibly bound electron transmitter plastocyanin, and they allocate the correct identity and orientation to all pigments, as well as to 621 [water molecules](#) that influence the energy transmission pathways. In connection with the loss of a second gene (*pgr5*), the genetically induced down-regulation of the

subunit Lhca2 results in the very efficient production of hydrogen in the double mutant.

Michael Hippler says, "The depletion of Lhca2 promotes the formation of PSI dimer, and so we suggest that the hydrogenase may favor the targeting of photosynthetic electrons from the PSI dimer, as we proposed in our earlier work. The structure of the PSI dimer enables us to make targeted genetic modifications in order to test the hypothesis of improved hydrogen production through the PSI dimer."

More information: Andreas Naschberger et al, Algal photosystem I dimer and high-resolution model of PSI-plastocyanin complex, *Nature Plants* (2022). [DOI: 10.1038/s41477-022-01253-4](https://doi.org/10.1038/s41477-022-01253-4)

Provided by University of Münster

Citation: Researchers provide new insights into photosynthesis (2022, October 14) retrieved 25 June 2024 from <https://phys.org/news/2022-10-insights-photosynthesis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.