

October 6 2022

# Achieving greater entanglement: Milestones on the path to useful quantum technologies



(a) Operating principle of our approach. Bell pairs are generated sequentially. The detection of one photon triggers the feed-forward including a field programmable gate array (FPGA), which in turn controls the operation mode of an all-optical storage loop. Possible operation modes are "read in and read out" (orange), "Storage" (green), or "PBS interference" (purple) selected by an appropriate switching of the electro-optic modulator (EOM). 2*N*-fold coincidences confirm the buildup of 2*N*-photon GHZ states. (b) Sketch of the experimental setup. A Ti:sapphire laser with a wavelength of 775 nm pumps a polarization Bell-state source based on parametric down-conversion in a Sagnac configuration (blue area). One photon of each emitted Bell pair is detected and triggers the feed-forward (red arrows), and the other photon is sent to our all-optical storage loop (green area), where it is stored until it is brought to



interference with the subsequent qubit. Credit: *Physical Review Letters* (2022). DOI: 10.1103/PhysRevLett.129.150501

Tiny particles are interconnected despite sometimes being thousands of kilometers apart—Albert Einstein called this "spooky action at a distance." Something that would be inexplicable by the laws of classical physics is a fundamental part of quantum physics. Entanglement like this can occur between multiple quantum particles, meaning that certain properties of the particles are intimately linked with each other.

Entangled systems containing multiple <u>quantum particles</u> offer significant benefits in implementing quantum algorithms, which have the potential to be used in communications, <u>data security</u> or quantum computing. Researchers from Paderborn University have been working with colleagues from Ulm University to develop the first programmable optical quantum memory. The study was published as an "Editor's suggestion" in the *Physical Review Letters* journal.

## **Entangled light particles**

The Integrated Quantum Optics group led by Prof. Christine Silberhorn from the Department of Physics and Institute for Photonic Quantum Systems (PhoQS) at Paderborn University is using minuscule light particles, or <u>photons</u>, as <u>quantum systems</u>. The researchers are seeking to entangle as many as possible in large states. Working together with researchers from the Institute of Theoretical Physics at Ulm University, they have now presented a new approach.

Previously, attempts to entangle more than two particles only resulted in very inefficient <u>entanglement</u> generation. In some cases, if researchers wanted to link two particles with others, it involved a long wait, as the



interconnections that promote this entanglement only operate with limited probability rather than at the touch of a button. This meant that the photons were no longer a part of the experiment once the next suitable particle arrived—as storing qubit states represents a major experimental challenge.

## **Gradually achieving greater entanglement**

"We have now developed a programmable, optical, buffer quantum memory that can switch dynamically back and forth between different modes—storage mode, interference mode and the final release," Silberhorn explains.

In the experimental setup, a small <u>quantum state</u> can be stored until another state is generated, and then the two can be entangled. This enables a large, entangled quantum state to grow particle by particle. Silberhorn's team has already used this method to entangle six particles, making it much more efficient than any previous experiments. By comparison, the largest ever entanglement of photon pairs, performed by Chinese researchers, consisted of twelve individual particles. However, creating this state took significantly more time, by orders of magnitude.

The quantum physicist explains: "Our system allows entangled states of increasing size to be gradually built up—which is much more reliable, faster, and more efficient than any previous method. For us, this represents a milestone that puts us in striking distance of practical applications of large, entangled states for useful quantum technologies." The new approach can be combined with all common photon-pair sources, meaning that other scientists will also be able to use the method.

**More information:** Evan Meyer-Scott et al, Scalable Generation of Multiphoton Entangled States by Active Feed-Forward and Multiplexing, *Physical Review Letters* (2022). DOI:



### 10.1103/PhysRevLett.129.150501

#### Provided by Paderborn University

Citation: Achieving greater entanglement: Milestones on the path to useful quantum technologies (2022, October 6) retrieved 27 June 2024 from <u>https://phys.org/news/2022-10-milestones-path-quantum-technologies.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.