


Scientists count electric charges in a single catalyst nanoparticle down to the electron

October 13 2022



Ultrahigh sensitivity and precision electron holography measurements around a platinum nanoparticle like the one shown here have allowed scientists to count the net charge in a single catalyst nanoparticle with a precision of just one electron for the first time. Credit: Murakami Lab, Kyushu University

If you often find yourself off by one when counting your socks after

doing the laundry, you might want to sit down for this.


Scientists in Japan have now counted the number of extra—or missing—[charges](#) down to a precision of just one electron in single platinum nanoparticles having diameters only one-tenth those of common viruses.

This new process for precisely studying differences in net charge on metal nanoparticles will aid in the further understanding and development of catalysts for breaking down greenhouse and other harmful gases into fuels and benign gases or for efficiently producing ammonia needed for fertilizers used in agriculture.

Led by Kyushu University and Hitachi Ltd., the research team achieved this feat of extreme counting through hardware and software improvements that increased tenfold the sensitivity of a technique called electron holography.

While [transmission electron microscopy](#) uses a beam of electrons to observe materials down to the [atomic level](#), [electron holography](#) utilizes the wave-like properties of electrons to probe electric and magnetic fields.

Interaction of an electron with fields causes a [phase shift](#) in its wave that can be identified by comparing it with a reference wave of an unaffected electron.



This new study highlights the importance of directly counting electric charges in a catalyst nanoparticle. For example, in a platinum nanoparticle on a surface of titanium oxide, the visualization of potential distribution by the developed noise reduction process in electron holography revealed negative charging of the nanoparticle with just six extra electrons. This is the first time charges per catalyst nanoparticle were counted with an accuracy of one electron charge. Credit: Murakami Lab, Kyushu University

In the new work, the researchers focused their microscopes on single nanoparticles of platinum on a surface of titanium oxide, a combination of materials that is already known to act as a catalyst and speed up chemical reactions.

On average, the platinum nanoparticles had diameters of only 10 nm—so small that it would take nearly 100,000 to span one millimeter.

"While each particle contains a few tens of thousands of atoms of platinum, the addition or removal of just one or two negatively charged electrons causes significant changes in the behavior of the materials as

catalysts," says Ryotaro Aso, associate professor at Kyushu University's Faculty of Engineering and first author on the paper in the journal *Science* reporting the work.

Measuring the fields just around a platinum nanoparticle—which vary depending on the imbalance of positive and [negative charges](#) in the particle—in an environment free of air, the researchers could determine the number of extra or missing electrons that are creating the fields.

"Amongst the millions of positively charged protons and negatively charged electrons balancing each other out in the nanoparticle, we could successfully tell if the number of protons and electrons was different by just one," explains Aso.

Although the fields are too weak to observe with previous methods, the researchers improved sensitivity by using a state-of-the-art 1.2-MV atomic-resolution holography microscope developed and operated by Hitachi that reduces mechanical and electrical noise and then processing the data to further tease out the signal from the noise.



Since 1966, Hitachi has been developing the holography electron microscope as an instrument for the direct observation of electric and magnetic fields in extremely small regions, and in 2014, developed a 1.2-MV atomic-resolution holography electron microscope with a grant under the Funding Program for World-Leading Innovative R&D on Science and Technology (the “FIRST Program”), a national project sponsored by the Japanese government. Credit: Hitachi, Ltd.

Developed by Osaka University's Yoshihiro Midoh, one of the paper's co-authors, the [signal processing](#) technique utilized the so-called wavelet hidden Markov model (WHMM) to reduce the noise without also removing the extremely weak signals of interest.

In addition to identifying the charge state of individual nanoparticles, the

researchers were able to relate differences in the number of electrons, which ranged from one to six, to differences in the crystal structure of the [nanoparticles](#).

While the number of electrons per area has been previously reported by averaging over a large-area measurement of many particles, this is the first time scientists could measure a single electron difference in a single particle.

"By combining breakthroughs in microscopy hardware and signal processing, we are able to study phenomenon on increasingly smaller levels," comments Yasukazu Murakami, professor at Kyushu University's Faculty of Engineering and supervisor of the Kyushu U team.

"In this first demonstration, we measured the charge on a single nanoparticle in vacuum. In the future, we hope to overcome the challenges that currently prevent us from doing the same measurements in the presence of gas to get information in environments closer to actual applications."

More information: Ryotaro Aso et al, Direct identification of the charge state in a single platinum nanoparticle on titanium oxide, *Science* (2022). [DOI: 10.1126/science.abq5868](https://doi.org/10.1126/science.abq5868).
www.science.org/doi/10.1126/science.abq5868

Provided by Kyushu University

Citation: Scientists count electric charges in a single catalyst nanoparticle down to the electron (2022, October 13) retrieved 26 June 2024 from <https://phys.org/news/2022-10-scientists-electric-catalyst-nanoparticle-electron.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.