


SWIFT J0503.7-2819 is an intermediate polar, research suggests

October 3 2022, by Tomasz Nowakowski



TESS light curve of J0503. Inset of the figure shows a close-up of some variability cycles. Credit: Rawat et al., 2022.

Using data from various space telescopes, astronomers have investigated a variable X-ray source known as SWIFT J0503.7-2819. Results of the study suggest that this source is an intermediate polar. The finding was detailed in a paper published September 22 on the arXiv pre-print server.

Cataclysmic variables (CVs) are [binary star systems](#) consisting of a white

dwarf accreting material from a normal star companion. They irregularly increase in brightness by a large factor, then drop back down to a quiescent state. Polars are a subclass of cataclysmic variables distinguished from other CVs by the presence of a very strong magnetic field in their [white dwarfs](#).

In some CVs, accretion occurs through a truncated accretion disk when the white dwarf is moderately magnetic. These systems are known as intermediate polars (IPs). Observations have shown that in IPs, the magnetic white dwarf spins asynchronously with the orbital period of the system and therefore produces a rapid oscillation with the spin period. Therefore, determining precise spin period and precise oscillation ephemeris could be the key to reveal the IP nature of some CVs.

At a distance of some 2,730 [light years](#), SWIFT J0503.7-2819 (or J0503 for short) is a variable X-ray source first identified in 2015. Further observations of J0503 have found that it has an orbital period of 81.6 minutes and that it may be an IP.

Therefore, in order to verify this hypothesis, a team of astronomers led by Nikita Rawat of the Aryabhata Research Institute of Observational sciences (ARIES) in India, performed X-ray observations of J0503 with NASA's Swift and ESA's XMM-Newton telescopes, and optical observations using NASA's Transiting Exoplanet Survey Satellite (TESS). The study was complemented by data from the American Association of Variable Star Observers (AAVSO) catalog.

"We have carried out X-ray and optical timing analyses and X-ray spectral analysis of a probable IP J0503," the researchers wrote in the paper.

The observations found that J0503 has an [orbital period](#) of approximately 81.65 minutes. Moreover, a 65.5-minute periodicity was

detected and the astronomers attribute it to the spin period of the white dwarf. If this scenario is true, then J0503 would be the first nearly synchronous IP below the period gap.

The [astronomers](#) suppose that J0503 is not synchronized due to the fact that the secondary star has a low magnetic moment. Therefore, it is unable to come into synchronism, similar to the nearly synchronous IPs and EX Hya-like systems.

The research found that J0503 might be accreting predominantly via stream-fed accretion. Furthermore, the identified energy-dependent spin pulsations appear to be due to the photoelectric absorption in the accretion flow, which is one of the unique characteristics of the majority of the IPs.

The study also derived fundamental parameters of the white dwarf in J0503. According to the paper, this star has a mass of about 0.54 solar masses, mass [accretion](#) rate at a level of 1.44×10^{-10} [solar masses](#) per year, and shock height of approximately 27.6 kilometers.

More information: Nikita Rawat, J. C. Pandey, Arti Joshi, Simone Scaringi, Umesh Yadava, SWIFT J0503.7-2819: A nearly synchronous intermediate polar below the period gap? arXiv:2209.11141v1 [astro-ph.HE], arxiv.org/abs/2209.11141

© 2022 Science X Network

Citation: SWIFT J0503.7-2819 is an intermediate polar, research suggests (2022, October 3) retrieved 6 February 2023 from <https://phys.org/news/2022-10-swift-j05037-intermediate-polar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.