

Planting and management of improved grass in tropical savannas can increase soil carbon by 15%

November 10 2022



Scientists tested Urochloa humidicola on an 8,000-hectare cattle ranch in Vichada, Colombia, and found soil carbon stocks 15 per cent above the control grassland. Credit: Hacienda San José



An improved variety of grass combined with a rotational grazing system can boost levels of soil carbon on tropical savannas by 15% compared to degraded pasture, a new study performed by the Alliance of Bioversity International and the International Center for Tropical Agriculture (CIAT), the University of Stuttgart, and the World Bank has found.

Scientists tested Urochloa humidicola, or Koronivia grass, on an 8,000-hectare cattle ranch in Vichada, Colombia, and found soil carbon stocks 15% above the control grassland, which itself had 40% higher carbon stocks than the default value provided by the IPCC.

The Koronivia grass captures carbon from the atmosphere through photosynthesis and enhance its storagein deep soil layers through its root systems.

The results, published in *Frontiers in Climate*, also found the use of improved grass on the natural savanna reduced <u>nitrous oxide</u> (N₂O) <u>emissions</u> from grazing cattle by a factor of 10, offsetting the emissions involved in cattle production. N₂O is a more potent greenhouse gas than <u>carbon dioxide</u>.

"This study shows, for the first time, the extent to which improved forages can help achieve a negative carbon footprint for livestock production," said Jacobo Arango, co-author of the paper and environmental biologist at the Alliance of Bioversity International and CIAT.

Dr. Arango, who was also a lead author on the latest IPCC report on mitigation, added, "At this point, reducing emissions is not enough; we must actively sequester carbon from the atmosphere. In our study, Koronivia grass achieved both, offering a way to protect the environment as well as food and livelihoods generated by livestock."



Cattle are responsible for roughly 9.5% of all anthropogenic greenhouse gas emissions worldwide. This includes <u>nitrous oxide emissions</u>, which are roughly 300 times more potent than carbon dioxide at heating Earth's atmosphere.

While <u>carbon sequestration</u> and lower greenhouse gas emissions are critical to meeting the Paris Agreement targets, few feasible solutions have been proven to work until now. Reducing nitrous oxide emissions in particular has been highlighted as a priority by President Biden ahead of the UN COP27 climate talks in Egypt this month.

The Koronivia grass was sourced from germplasm at the Future Seeds gene bank, which contains the world's largest collection of tropical forages and is managed by CGIAR's Alliance of Bioversity International and the International Center for Tropical Agriculture (CIAT). The facility, which is based near Cali, Colombia, helps to preserve crucial plant biodiversity while supporting cutting-edge agricultural research.

As well as reducing emissions and boosting <u>soil carbon</u> storage, the grass can also grow on marginal land, meaning farmers can make more productive use of degraded soils without resorting to converting virgin land.

"This research represents a great gift from Colombia to the world. It's a way to act concretely in favor of global warming mitigation and the promotion of food safety and the well-being of thousands of cattle farmers," stated Gabriel Jaramillo, from Hacienda San José, where the study was conducted.

"At Hacienda San Jose we are approaching sustainability with a holistic view. This allows us to work considering human and animal well-being, harmony with the environment, virgin forests and wetlands preservation, and the implementation of best practices like the ban of pre-emptive



antibiotics and the annual burning of savannas. Working with and educating the community is also crucial to improve their living standards," Jaramillo concluded.

More information: Ciniro Costa et al, Soil carbon stocks and nitrous oxide emissions of pasture systems in Orinoquía region of Colombia: Potential for developing land-based greenhouse gas removal projects, *Frontiers in Climate* (2022). DOI: 10.3389/fclim.2022.916068

Provided by The Alliance of Bioversity International and the International Center for Tropical Agriculture

Citation: Planting and management of improved grass in tropical savannas can increase soil carbon by 15% (2022, November 10) retrieved 26 April 2024 from <u>https://phys.org/news/2022-11-grass-tropical-savannas-soil-carbon.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.