

International team observes innermost structure of quasar jet

GMVA+ALMA 3 mmHSA 2 cmHST OpticalHSA
x20 zoomHSA
x2160 zoomHST
x2160 zoomHST
x2160 zoomI Ight years
20 microarcsecondsI Ight years
x20 microarcsecondsI Ight years
x20 microarcsecondsI Ight years
x20 microarcseconds

Radio astronomy images of the 3C 273 jet. The close-up view on the left is the deepest look yet into the plasma jet of the quasar 3C 273. The image in the center shows the extended structure of the jet. The image on the right is a visible light image of the quasar taken by the Hubble Space Telescope. The radio observations were made by the Global Millimeter VLBI Array (GMVA) joined by the Atacama Large Millimeter/submillimeter Array (ALMA) and the High Sensitivity Array (HSA). Credit: Hiroki Okino and Kazunori Akiyama; GMVA+ALMA and HSA images: Okino et al.; HST Image: ESA/Hubble & NASA

November 22 2022



An international team of scientists has observed the narrowing of a quasar jet for the first time by using a network of radio telescopes across the world. The results suggest that the narrowing of the jet is independent of the activity level of the galaxy which launched it.

Nearly every galaxy hosts a <u>supermassive black hole</u> in its center. In some cases, enormous amounts of energy are released by gas falling towards the black hole, creating a phenomenon known as a quasar. Quasars emit narrow, collimated jets of material at nearly the speed of light. But how and where quasar jets are collimated has been a longstanding mystery.

An international team led by Hiroki Okino, a graduate student at the University of Tokyo, and including members from the National Astronomical Observatory of Japan (NAOJ), the Massachusetts Institute of Technology, Kogakuin University, Hachinohe National College of Technology, and Niigata University, captured an image with the highest angular resolution to date that shows the deepest part of the jet in a bright quasar known as 3C 273.



Multifrequency images of 3C 273 jets with the HSA at 43, 22, and 15 GHz and the VLBA at 1.7 GHz. Each panel shows the mean top-set image at each frequency restored by the circular Gaussian beam with the beam solid angle of the uniform-weighting beam size in Table <u>1</u>. The lowest contour levels are 6.5



mJy beam⁻¹ at 43 GHz, 8.2 mJy beam⁻¹ at 22 GHz, 7.1 mJy beam⁻¹ at 15 GHz, and 11.6 mJy beam⁻¹ at 1.7 GHz. The contours for all images are multiplied by a factor of 2. The lowest contour level of each image is estimated from the mean residual image rms of the top-set reconstructions. Credit: *The Astrophysical Journal* (2022). DOI: 10.3847/1538-4357/ac97e5

The team found that the jet flowing from the quasar narrows down over a very long distance. This narrowing part of the jet continues incredibly far, well beyond the area where the black hole's gravity dominates. The results show that the structure of the jet is similar to jets launched from nearby <u>galaxies</u> with a low luminosity active nucleus. This would indicate that the collimation of the jet is independent of the activity level in the host galaxy, providing an important clue to unraveling the inner workings of jets.

These results appeared in *The Astrophysical Journal* on November 22, 2022.

More information: Hiroki Okino et al, Collimation of the Relativistic Jet in the Quasar 3C 273, *The Astrophysical Journal* (2022). DOI: 10.3847/1538-4357/ac97e5

Provided by National Astronomical Observatory of Japan

Citation: International team observes innermost structure of quasar jet (2022, November 22) retrieved 1 May 2024 from https://phys.org/news/2022-11-international-team-innermost-quasar-jet.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.