

# **Dam safety: Study indicates probable maximum flood events will significantly increase over next 80 years**

November 15 2022, by Neil Martin

---



Credit: AI-generated image ([disclaimer](#))

The rainfall model that engineers use to help them design critical infrastructure such as large dams and nuclear power plants needs to be updated to take into account climate change.

That is the conclusion of a paper published in *Water Resources Research*, which calculates that the "Probable Maximum Precipitation" (PMP) estimates for 546 [large dams](#) across Australia are expected to increase between 14 and 38% on average due to increasing atmospheric moisture.

The research suggests that existing dams will be at greater risk under [climate change](#) than what is currently assumed.

The researchers say existing models of PMP have not been updated for at least 20 years, and more recent meteorological events already show that the climate is warming and making storms more intense and more frequent.

"There are a lot of risks involved with dams given the amount of water they are holding back. Some of the worst floods around the world were due to extreme storms overwhelming a dam, causing it to fail and release a wall of water downstream," says Johan Visser from UNSW Sydney's Water Research Center, who was lead author on the paper in collaboration with academics from the University of Melbourne.

"Engineers design dams to accommodate the largest flood event that could reasonably be expected to occur at a particular location, known as the Probable Maximum Flood—PMF. To work that out, you first need to calculate what is the greatest depth of rainfall meteorologically possible over that area in a certain amount of time, which we call the Probable Maximum Precipitation—PMP.

"The problem is that PMP calculation is based solely on [historical data](#) with no consideration for future climate conditions. This means that many large dams constructed decades ago were designed using information representative of a cooler climate.

"The purpose of this research was to analyze whether PMP estimates

have changed over the last six decades and how these estimates might change in the future if we take into consideration a potential increase in atmospheric moisture due to known climate change," says Mr Visser.

## **Incorporation of climate models**

The current PMP guidelines for various timeframes and locations across Australia are collated and published by the Bureau of Meteorology.


The new research—which was funded by 10 of Australia's leading water providers and dam owners from across Australia—reanalyzed existing meteorological records and added in more up-to-date data from the last 20 years that was not previously included. The researchers then calculated potential changes in the future by incorporating the latest climate scenario modeling from the highly respected Coupled Model Intercomparison Project Phase 6 (CMIP6).

These [climate models](#) are used to explore how a range of global socioeconomic choices over the next century will affect greenhouse gas emissions.

The results of the paper show that PMP estimates have increased across Australia over the past 60 years and are expected to increase further due to predicted increases in atmospheric moisture. Based on the trajectory of the observed data, it was evident there would be a systematic increase in the PMP. This was confirmed using climate model simulations, indicating further increases for every climate scenario analyzed.

For the worst-case future scenario, where green policies are not implemented and carbon emissions remain highest through to the year 2100, PMP estimates over large dam locations in Australia could increase by 38% on average.

Even using the most conservative (low) estimates regarding emissions and subsequent climate change, the modeling suggests an average increase in PMP of 13% across Australia.



Results of the study show that maximum persisting dewpoints (input in PMP calculations) over Australia has increased over the historical record and is expected to increase further over the next 80 years for all modelled climate scenarios known as Shared Socioeconomic Pathways (SSPs), resulting in increased PMP estimates. Credit: University of New South Wales

The researchers say they have shown the current method of calculating PMP is likely outdated and does not take into consideration the potential consequences of current changes in atmospheric conditions, let alone

those predicted into the future.

"For each 1-degree Celsius rise in temperature, the atmosphere can hold approximately 7% more water," says Mr. Visser.

"We can already see in the observational record that extreme rainfall events and floods are becoming more frequent and more intense.

"The observational data shows that atmospheric moisture has been increasing over the historical record, and when you also factor in future climate projections from climate models, then you see that it's projected to increase even further into the future.

"That is true for all the scenarios we analyzed, even the most conservatively low estimates where there's an increasing shift toward sustainable practices. So, under any future scenario, our estimate is that the PMP will increase.

"That is concerning because if PMPs are increasing, then our PMF estimates will also increase. Dam owners regularly review the safety performance of their dams under a wide range of failure scenarios, and this research reinforces the importance of ensuring that in the future these reviews consider the implications of a warming climate on extreme floods.

"One of the issues with the methodology presently used to estimate probable maximum rainfalls is that the last time it was updated in Australia was around the late 1990s or early 2000s. Therefore, 20 years' worth of atmospheric moisture and rainfall data is already not included."

Professor Rory Nathan of the University of Melbourne, one of the lead collaborators involved in the research, said, "No country in the world has yet updated the operational procedures used to estimate PMPs to account



for climate change, and this research provides the first evidence that these procedures need to be reviewed.

"It will take some time to work out the best way of using these findings to update industry practice, but we are facing a climate crisis and this research adds to the urgency with which we need to curb our carbon emissions."

## **Global importance of PMP modeling**

The researchers say it is hard to make a generalized recommendation with regards to potential upgrades to dam infrastructure across Australia based on the implications of the results of their new study.

That's because the location, climate and design of each dam varies significantly, and individual risk and cost assessments would need to be done on a case-by-case basis. It also needs to be stressed that climate change is continually evolving, and the increase in extreme rainfalls associated with a warming climate is just one of the factors that dam owners need to consider when regularly reviewing the safety performance of their dams.

But they highlight the fact that previous amendments to PMP estimates have resulted in large-scale dam enhancements.

For example, the Warragamba Dam that supplies water to Sydney had its wall raised by 5.1 meters and an auxiliary spillway (18,000 m<sup>3</sup>/s capacity) constructed after hydrological studies in the 1980s showed the original spillway (10,000 m<sup>3</sup>/s capacity) was undersized and the risk of dam break was unacceptably high.

Professor Ashish Sharma, co-author of the paper from UNSW's School of Civil and Environmental Engineering, said, "This is an important

piece of research, not just for Australia but also across the world. The approach to PMP is developed and recommended by the World Meteorological Organization and is used more or less consistently by countries around the globe.

"Anybody who is planning to build a large dam needs to think 50 to 100 years ahead, and this research makes clear that future planning now needs to consider the future impacts of climate change on the plausible upper limit of rainfall extremes."

**More information:** J.B. Visser et al, The impact of climate change on operational estimates of Probable Maximum Precipitation, *Water Resources Research* (2022). [DOI: 10.1029/2022WR032247](https://doi.org/10.1029/2022WR032247)

Provided by University of New South Wales

Citation: Dam safety: Study indicates probable maximum flood events will significantly increase over next 80 years (2022, November 15) retrieved 20 March 2024 from <https://phys.org/news/2022-11-safety-probable-maximum-events-significantly.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--