

Mapping street-level pollution estimates to reveal safer routes

November 29 2022, by Morgan Rehnberg



Hong Kong is seen by satellite here in January 2018. Credit: <u>Axelspace</u> <u>Corporation/Wikimedia Commons</u>, <u>CC BY-SA 4.0</u>

The United Nations has identified improving the walkability and bikeability of cities as a key goal in efforts to reduce dependence on greenhouse gas-emitting automobiles and promote routine exercise for public health. However, increased walking and cycling can come with their own health risks.



Time spent in the open air, in addition to other factors, increases exposure to airborne pollutants, for example. The most harmful category of pollutants to human health includes inhalable particles smaller than 2.5 micrometers, referred to as $PM_{2.5}$. It has been estimated that more than 4 million <u>premature deaths</u> worldwide are attributable to these pollutants each year.

To help reduce health risks from this pollution, Tong et al. developed a new approach for estimating ground-level $PM_{2.5}$ concentrations from satellite observations and incorporated it into a mobile mapping application that identifies routes through Hong Kong to minimize pedestrians' exposure to $PM_{2.5}$.

The researchers combined several <u>data sources</u> to build their framework, which accounts for city-specific factors that produce and concentrate $PM_{2.5}$. Sixteen air quality monitoring stations provided hourly $PM_{2.5}$ estimates across the city. Landsat 8–derived data products provided maps of vegetation, impervious surfaces, and aerosol density. Weather data were incorporated through the use of NASA's Goddard Earth Observing System-Forward Processing (GEOS-FP) forecasts, and road data from OpenStreetMap helped quantify the contributions of automobile and bus traffic.





Two routes for pedestrians or bicyclists to take between Hong Kong Polytechnic University and Hong Kong Metropolitan University are shown. (a) The "healthy" route is 100 meters longer than (b) the shortest route, but $PM_{2.5}$ pollution is reduced by about 5%–10%. Credit: <u>Tong et al.</u>, <u>CC BY-NC-ND 4.0</u>

The input data were used to estimate real-time $PM_{2.5}$ concentrations on a grid across the city. That grid was then overlaid by a graph with nodes representing street intersections and edges representing the streets themselves. Each edge was assigned a pollution "cost" based on the $PM_{2.5}$ in the grid cells it crosses. The pollution breathed during a journey



between two nodes is the sum of the costs of the edges that most efficiently connect them.

The Hong Kong case study included $PM_{2.5}$ estimates for 70,788 roads. Using the developed app, which is visually similar to commercial mapping apps, a user can compare the fastest and healthiest routes between two points. The researchers observed that healthy route planning could reduce $PM_{2.5}$ exposure by 5%–25%. In one highlighted example, the app indicated that choosing a route between two Hong Kong universities that was 100 meters longer than the shortest available route would result in a 5%–10% reduction in breathed pollution.

The researchers note that the generalizability of their approach for estimating high-resolution $PM_{2.5}$ levels means it may be applicable in other cities and useful in other route planning applications and services to minimize people's pollution exposure.

The research is published in the journal GeoHealth.

More information: Chengzhuo Tong et al, Estimation of On-Road PM 2.5 Distributions by Combining Satellite Top-of-Atmosphere With Microscale Geographic Predictors for Healthy Route Planning, *GeoHealth* (2022). DOI: 10.1029/2022GH000669

This story is republished courtesy of Eos, hosted by the American Geophysical Union. Read the original story <u>here</u>.

Provided by Eos

Citation: Mapping street-level pollution estimates to reveal safer routes (2022, November 29) retrieved 26 April 2024 from <u>https://phys.org/news/2022-11-street-level-pollution-reveal-safer-routes.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.