


Leading the way in superconductor research: New compounds of lanthanum and hydrogen

November 22 2022, by Christian Wißler



Arrangement of lanthanum atoms in the lanthanum hydrides synthesized in this work. **a** LaH_3 ; **b** $\text{LaH}_{\sim 4}$; **c** $\text{LaH}_{4+\delta}$; **d** La_4H_{23} ; **e** $\text{LaH}_{6+\delta}$; **f** $\text{LaH}_{9+\delta}$; **g** $\text{LaH}_{10+\delta}$. The red spheres represent lanthanum atoms. Credit: *Nature Communications* (2022). DOI: 10.1038/s41467-022-34755-y

Superconducting materials are characterized by the fact that they lose their electrical resistance below a certain temperature, the so-called transition temperature. In principle, they would be ideal for transporting electrical energy over very long distances from the electricity producer to the consumer.

Numerous energy challenges would be solved in one fell swoop: For example, the electricity generated by [wind turbines](#) on the coast could be channeled inland without losses. However, this would only be possible if materials were available that have superconducting properties at normal room and ambient temperatures.

In 2019, an unusually high transition temperature of minus 23 degrees Celsius was measured in experiments coordinated by the Max Planck Institute in Mainz. The measurement took place at a compression pressure of 170 gigapascals—1.7 million times higher than the pressure of the Earth's atmosphere. The material was a lanthanum [hydride](#) ($\text{LaH}_{10+\delta}$), a compound of atoms of the metal lanthanum with [hydrogen](#) atoms. The report on these experiments and other similar reports remain highly controversial. They have internationally aroused great interest in research on lanthanum hydrides with different compositions and structures.

The new study, published in *Nature Communications*, takes up this focus of research. The measurement data from 2019 suggested that also other superconducting lanthanum hydrides form under very high compression pressures.

These considerations have now been confirmed: A total of seven lanthanum hydrides were produced in the high-pressure laboratory of the Bavarian Research Institute of Experimental Geochemistry & Geophysics (BGI): the two already known compounds $\text{LaH}_{10+\delta}$ and LaH_3 , and the previously unknown lanthanum hydrides $\text{LaH}_{\sim 4}$, $\text{LaH}_{4+\delta}$, La_4H_{23} , $\text{LaH}_{6+\delta}$, and $\text{LaH}_{9+\delta}$.

All these compounds were formed from samples containing lanthanum and paraffin which is a hydrogen-rich mixture of saturated hydrocarbon. The samples were subjected to very high pressures between 96 and 176 gigapascals in diamond anvil cells and heated to over 2,200 degrees

Celsius.

In cooperation with the German Electron Synchrotron (DESY) in Hamburg and the Center for Advanced Radiation Sources in Chicago, it was possible to identify the structures of the new compounds of lanthanum and hydrogen. It turned out that lanthanum hydrides with the same arrangement of lanthanum atoms differ considerably in their hydrogen content.

In other words, the same framework of lanthanum atoms can be linked to different numbers of hydrogen atoms. The hydrogen atoms can thereby be arranged in very different ways. The scientists have shown that a similar structural diversity can also exist in hydrides that contain other metals from the rare earth group instead of lanthanum.

These surprising findings refute a hypothesis that has played a central role in research into [superconducting materials](#) up to now: namely, the prejudice that a certain number and arrangement of lanthanum atoms only allows only for one specific configuration of hydrogen atoms.

Against this background, the coordinator of the study, Prof. Dr. Dr. h.c. Natalia Dubrovinskaia from the Laboratory of Crystallography at the University of Bayreuth explains, "In our search for superconductors with higher transition temperatures, [theoretical models](#) and calculations based on them are indispensable. Hydrogen-containing solids have proven to be highly promising materials."

"The superconductivity of these chemical compounds depends, as we know today, essentially on the number and arrangement of the hydrogen atoms. It is therefore all the more important that our theoretical models do not incorporate incorrect assumptions that lead to hydrogen-containing solids with a high transition temperature remaining undiscovered."

Prof. Dr. Dr. h.c. Leonid Dubrovinsky from BGI adds, "Our findings on [lanthanum](#) hydrides firmly remind us that in the search for optimal superconductors we must not underestimate the number of possible hydrogen-containing compounds and the variety of possible configurations of [hydrogen atoms](#)."

More information: Dominique Laniel et al, High-pressure synthesis of seven lanthanum hydrides with a significant variability of hydrogen content, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-34755-y](#)

Provided by Bayreuth University

Citation: Leading the way in superconductor research: New compounds of lanthanum and hydrogen (2022, November 22) retrieved 3 May 2024 from <https://phys.org/news/2022-11-superconductor-compounds-lanthanum-hydrogen.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--