

## Sweet corn sweltering in summer heat spells uncertainty for corn lovers





Nonlinear relation between temperature and yield for the (A) Midwest-Rainfed, (B) Midwest-Irrigated, and (C) Northwest-Irrigated. Graphs at the top of each frame display changes in log yield if the crop is exposed for one day (24 h) to a particular 1 °C temperature interval. Exposure (days) was calculated as the sum of the fraction of a day during which temperatures fall within each interval. The 95% confidence intervals are shown by gray shaded area. Histograms at the bottom of each frame display the average temperature exposure for all yield observations in each production region. Credit: *Scientific Reports* (2022). DOI: 10.1038/s41598-022-23237-2

Few things say summer in America more than buttery corn on the cob, but as summer temperatures climb to unprecedented levels, the future of sweet corn may not be so sweet. New University of Illinois research shows sweet corn yields drop significantly with extreme heat during



flowering, especially in rainfed fields in the Midwest.

Climate projections don't just predict a handful of hot days going forward. The U.S. Global Change Research Program predicts 20 to 30 more days over 32 C [about 90 F] by mid-century across much of the U.S.

"The reality is that producing sweet corn, one of the most popular vegetable crops in the U.S., will be more difficult in the future. We need to develop new approaches and technologies to help crops adapt to <u>climate change</u>," says Daljeet Dhaliwal, former graduate research assistant and lead author on the study published in *Scientific Reports*.

Dhaliwal worked with Marty Williams, USDA-ARS ecologist and affiliate professor in the Department of Crop Sciences at Illinois, to document yield response of sweet corn to growing season temperature and precipitation over a 27-year period. Williams obtained private data from sweet corn processors for 16,040 individual fields in Illinois, Minnesota, Washington, and Wisconsin, providing a much finer resolution than similar studies in field corn using county-level data.

"Our analysis reveals that small temperature changes have a greater influence on crop yield compared to small precipitation changes for both rainfed and irrigated fields in the Midwest and Northwest, but rainfed production shows greater sensitivities," Williams says.

He adds that <u>extreme temperatures</u> during flowering can influence pollen viability, fertilization, kernel abortion, and other processes.

"If there's a bad time for extreme heat, it's during flowering. That's especially true in a crop where ear quality is so important. With <u>heat</u> <u>stress</u> during flowering, you can have ears with fewer kernels or very misshapen kernels that look nothing like what the consumer is



expecting," Williams says.

This study used the concept of "extreme degree days" to capture the cumulative effect of temperatures above 30 C [86 F] during flowering on sweet corn yield. Degree days are normally calculated by taking the average of high and low temperatures over a given 24-hour period. To calculate extreme degree days, Dhaliwal summed degree days over 30 C.

For rainfed regions, every extreme degree day during flowering led to an additional yield loss of 2%. For irrigated systems, yield loss was less severe; just 0.5% per degree day over 30 C.

But more than one day at 40 C [104 F] could drop yield by 20%. This result was based on season-long hourly <u>temperature</u> data, not degree days.

The pattern suggests irrigation could ameliorate the effects of <u>extreme</u> <u>heat</u>, but climate scenarios also warn water could be in short supply in the future. The authors note that if sweet corn yield and quality continues to decline due to climate extremes, planting times or production areas may need to shift to avoid the hottest temperatures.

In the meantime, Williams says there's only so much breeders can do to build in heat tolerance.

"The question becomes how do we develop sweet <u>corn</u> to become more resilient to increasing environmental stressors? To what extent can you do that by managing the crop versus improving the crop itself? These are hard questions.

"The thing we really need to do to prevent worse outcomes is to minimize further climate change. But we're in a period where we're going have to deal with what's already set in motion," Williams says.



"Our trajectory is already set for the next several decades."

**More information:** Daljeet S. Dhaliwal et al, Evidence of sweet corn yield losses from rising temperatures, *Scientific Reports* (2022). DOI: 10.1038/s41598-022-23237-2

## Provided by University of Illinois at Urbana-Champaign

Citation: Sweet corn sweltering in summer heat spells uncertainty for corn lovers (2022, November 22) retrieved 3 May 2024 from <u>https://phys.org/news/2022-11-sweet-corn-sweltering-summer-uncertainty.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.