


A research agenda for nonvascular photoautotrophs under climate change

December 13 2022



Impacts of nonvascular photoautotrophs (NVP) on key aspects of ecosystem functioning (yellow boxes) in selected ecosystem types that show a high abundance of NVP. The effects are categorized according to the interconnected biogeochemical cycles of carbon (gray arrow), nutrients (mainly nitrogen and phosphorus) (orange arrow), and water and energy (blue arrow). The cycles are arranged according to the supposed primary impacts of NVP in the respective ecosystems. These impacts are, however, not limited to a specific ecosystem. Credit: *New Phytologist* (2022). DOI: 10.1111/nph.18631

Non-vascular photoautotrophs (NVP), including mosses, lichens, terrestrial algae and cyanobacteria, are organisms that receive their energy from light but lack the vascular tissue that transports water and nutrients in vascular plants. Lacking roots, they rely on direct uptake of water from the atmosphere or from the near-surface layer of the soil.

Because [water loss](#) cannot be actively controlled, the organisms often become dehydrated. However, unlike most [vascular plants](#), NVP are able to largely adapt their metabolism to these large fluctuations in water content, making them a common form of vegetation in many arid ecosystems, such as deserts, tundras, and high elevations.

NVP are considered essential for the functioning of ecosystems in many regions of the world since they are, for instance, responsible for about 50% of nitrogen input to [natural ecosystems](#) through their association with [nitrogen-fixing bacteria](#), control the partitioning as well as evapotranspiration of precipitation in forests, have an impact on near-ground air temperature, contribute to peatland ecosystems that store an estimated 30% of global soil carbon, or form biocrusts that protect soil surfaces from erosion by water and wind.

Current research suggests that climate change may pose a significant threat to NVP, with major impacts on many regions of the world. But the extent to which this will affect associated ecosystems is highly uncertain.

"We found that ecosystem functions of NVP are likely to be significantly affected by [climate change](#) and that a better quantitative understanding of some key processes is needed, for example, the potential for acclimation, the response to elevated carbon dioxide, the role of the microbiome, and the feedback from ecosystem changes to climate. We propose an [integrative approach](#) with innovative, cross-method laboratory and field experiments and ecophysiological modeling,

for which sustained scientific collaboration in NVP research is essential," reports the study's first author, Prof. Dr. Philipp Porada of the Department of Biology at the Universität Hamburg.

The paper is published in the journal *New Phytologist*.

More information: Philipp Porada et al, A research agenda for nonvascular photoautotrophs under climate change, *New Phytologist* (2022). [DOI: 10.1111/nph.18631](https://doi.org/10.1111/nph.18631)

Provided by University of Hamburg

Citation: A research agenda for nonvascular photoautotrophs under climate change (2022, December 13) retrieved 23 April 2024 from <https://phys.org/news/2022-12-agenda-nonvascular-photoautotrophs-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.