



# Drying process could be key step in development of life

December 21 2022, by Jason Daley

**(a) Mechanism 1**



**(b) Mechanism 2**



Credit: *Origins of Life and Evolution of Biospheres* (2022). DOI: 10.1007/s11084-022-09634-7

One-hundred fifty years ago, Charles Darwin speculated that life likely originated in a warm little pond. There, Darwin supposed, chemical reactions and the odd lightning strike might have led to chains of amino acids that, over time, became more and more complex until the beginnings of life emerged.

Ever since, researchers have investigated this type of pre-life or "prebiotic" chemistry, trying to figure out the [chemical](#) pathways that could have led from a pool filled with simple amino acids to bacteria, redwood trees and people. After a series of experiments, University of Wisconsin–Madison chemical engineering Ph.D. student Hayley Boigenzahn and John Yin, a professor of chemical and [biological engineering](#) and a founding faculty member of the Wisconsin Institute for Discovery, can explain how one of the potentially crucial early steps on the path of life could have happened. They published their findings in the journal *Origins of Life and Evolution of Biospheres*.

In a famous 1952 study called the Miller-Urey experiment, researchers simulated the conditions thought to be present on the prebiotic Earth, including certain ratios of water, methane, hydrogen and other elements. When zapped with electricity to simulate lightning, the researchers found that the reaction produced amino acids, suggesting that these molecules were widely present on the prebiotic Earth.

"We know amino acids are the building blocks of proteins and proteins are essential for life," says Yin. "In prebiotic chemistry, it's long been a question of how we could we get these things to form bonds and strings in a manner that might eventually lead to a living cell. The question is hard because the particular chemistry involved is one that tends to fail in the presence of water."

In her experiment, Boigenzahn investigated whether it's possible these amino acids could have come together during periods of environmental

change—for instance, as a pool of water evaporated. In the presence of a chemical activator, these amino acids could bond together into peptides, or short chains of amino acids.

To study how amino acids might form bonds during the drying process, Boigenzahn created solutions of the amino [acid](#) glycine and trimetaphosphate, an activator that is naturally created during volcanic processes. Using a heater to evaporate the solution, Boigenzahn watched what happened to the amino acids over 24 hours.

What she found was a two-stage process. In the first stage, when the pH of the solution was alkaline, the glycine combined into two-molecule units called dimers, which are also produced protons, making the pH of the solution neutral. In the second stage, as evaporation took place, the dimers began to bond together to form longer peptide chains, called oligoglycine.

It's easy to imagine a scenario in which amino acids in a volcanically warmed hot spring containing an activator first combine into dimers. Then, as the water evaporates and its chemistry changes, the dimers bond and begin to form into longer chains of [amino acids](#).

"What we're showing here is that that it doesn't necessarily have to be the same environment throughout all the reactions," says Boigenzahn. "They can occur in [different environments](#), provided that the reactions that are occurring help create an environment that's beneficial for the next steps."

Through multiple wet-dry cycles, it's possible the peptide chains grew longer and longer. Eventually, they could have begun to fold in on themselves, forming enzymes, or proteins that catalyze [chemical reactions](#). That could set the stage for more complex proteins and the beginnings of metabolism.

Boigenzahn and Yin both say it will be a long time before researchers figure out a possible path from Darwin's warm little pond to the beginnings of life. But, especially for [chemical engineers](#), the effort of studying prebiotic chemistry could have big payoffs.

"If you really understand this chemistry, which is different from traditional biology, eventually you might create chemical systems that are able to store information, adapt and evolve," says Yin. "DNA stores information at thousands of times the density of a computer chip can. If we could get systems that do this without necessarily being living cells, then you start to think about all sorts of new functions and processes occurring at the molecular level."

**More information:** Hayley Boigenzahn et al, Glycine to Oligoglycine via Sequential Trimetaphosphate Activation Steps in Drying Environments, *Origins of Life and Evolution of Biospheres* (2022). [DOI: 10.1007/s11084-022-09634-7](https://doi.org/10.1007/s11084-022-09634-7)

Provided by University of Wisconsin-Madison

Citation: Drying process could be key step in development of life (2022, December 21) retrieved 16 July 2024 from <https://phys.org/news/2022-12-drying-key-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.