


New method of reducing carbon dioxide could be a golden solution to pollution

December 8 2022



A gold-based hybrid material (Au@CB[6]) is modified by CB[6] in order to efficiently convert CO₂ to CO. Credit: *Nano Research*, Tsinghua University Press

Reducing carbon dioxide (CO₂) to carbon monoxide (CO) through an electrochemical reaction holds strong potential for removing CO₂ from

the atmosphere to reduce pollution and for producing alternative energy with carbon monoxide as an ingredient.

However, the current catalysts used in electrochemical [carbon dioxide](#) reduction reaction (CO₂RR) are not efficient or selective enough to make CO₂RR a practical solution. Now, a team of researchers from Fujian Institute of Research on the Structure of Matter of Chinese Academy of Sciences has developed a gold-based hybrid material by modifying [gold nanoparticles](#) with a macrocyclic compound called cucurbit[6]uril (CB[6]) that allows for more efficient CO₂RR than previously possible.

The results were published in *Nano Research*.

"With this work, we hoped to solve the problem of environmental pollution and energy shortage through electrochemical conversion of carbon dioxide to value-added products," said corresponding author Minna Cao of the State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, and of the University of Chinese Academy of Sciences.

"In order to enhance the local CO₂ concentration on the catalysts' surface, we utilize macromolecule cucurbit[n]uril to functionalize gold surface, which is the distinguishing feature of our work from those that has been done before."

According to the researchers, gold is known to be highly active in converting CO₂ to CO relative to other catalysts. However, the binding energy of both CO₂ and CO to the gold [catalyst](#) surface is positively correlated, which clashes with the need in CO₂RR for CO₂ adsorption and CO desorption, since the CO desorption does not occur due to the positive correlation of its [binding energy](#) to the catalyst.

The researchers created a controlled synthesis of nanoparticles by modifying CB[6]. CB[6] has negatively charged portals and positively charged surface, which helps contribute to the result of having the electronic interaction between CB[6] and metal regulate the catalytic performance.

The researchers verified both the morphology and surface structure of the nanoparticles through transmission electron microscopy. The gold-based hybrid material (Au@CB[6]) was proven to enhance CO₂RR catalytic activity.

"We have proved the interaction between cucurbit[6]uril and CO₂ through operando electrochemical measurement and density functional theory calculations," Cao said.

According to the researchers, multiple factors contribute to the improved catalytic performance. First, the CB[6] can increase the local CO₂ concentration near the [metal surface](#) by gathering CO₂, which means that the Au@CB[6] has tunable, or adjustable, CO₂ enrichment. Additionally, the modification of CB[6] allow for improved CO₂RR by breaking the previously mentioned scaling relations of the binding affinity between catalyst surface and CO₂/CO.

Also, one reason that CO₂RR was limited in efficiency with [gold surface](#) catalysts previously is that CO₂ has low solubility in aqueous electrolytes, an issue that the researchers solved by using the highly specific binding force of macrocycle to selectively adsorb certain species to regular the electrocatalytic reaction.

"The results showed that CB[6] can gather CO₂ and lead the increased local CO₂ concentration near the metal interface, as well as promote CO desorption, which are the dominating reasons for enhanced CO₂RR performance," Cao said. "Using the rigid macrocycle cucurbit[n]uril to

modify the catalysts' surface is a promising pathway to enhance the electrocatalytic performance."

The researchers plan to continue to modify the catalyst in order to further improve the efficiency of the CO₂RR.

"In the next step, we hope to adjust the shape and size of the gold catalyst in the presence of cucurbit[n]uril to further promote the catalytic performance toward electrochemical reduction of carbon dioxide to value-added products," Cao said.

More information: Huimin Wang et al, Tunable CO₂ enrichment on functionalized Au surface for enhanced CO₂ electroreduction, *Nano Research* (2022). [DOI: 10.1007/s12274-022-5159-8](https://doi.org/10.1007/s12274-022-5159-8)

Provided by Tsinghua University Press

Citation: New method of reducing carbon dioxide could be a golden solution to pollution (2022, December 8) retrieved 16 June 2024 from <https://phys.org/news/2022-12-method-carbon-dioxide-golden-solution.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.