

Researchers reveal structural evolution during molybdenum carbides formation at atomic level

December 1 2022, by Li Yuan



In situ investigations on structural evolutions during the facile synthesis of cubic α -MoC_{1-x} catalysts. Credit: Yu Jiafeng

Transition metal carbides have attracted much attention due to their unique catalytic properties similar to precious metals. Among them, α -



phase molybdenum carbides $(\alpha - MoC_{1-x})$ show high activity in activation of water at an ultra-low temperature in water-gas shift and aqueous-phase reforming of methanol reaction.

However, the preparation process of cubic α -MoC_{1-x} generally requires an extra ammonification treatment with high energy-consumption or assistance by a significant amount of <u>precious metals</u>, which severely limits its application.

Recently, a joint research team led by Prof. Sun Jian, Dr. Yu Jiafeng, and Dr. Liu Yuefeng from the Dalian Institute of Chemical Physics (DICP) of the Chinese Academy of Sciences (CAS), in collaboration with Prof. Grunwaldt from Karlsruhe Institute of Technology (KIT) of Germany, observed the dynamic evolution of <u>crystal structure</u> at the <u>atomic level</u>, which confirmed that the cubic MoO_x phase was the key intermediate for the synthesis of face-centered cubic α -MoC_{1-x} by onestep carburization process.

This work was published in *Journal of the American Chemical Society* on Nov. 23.

The researchers conducted qualitative and quantitative analyses of crystal phases formed as intermediates during the carburization process, and further monitored the evolution of morphology and crystal structure by environmental transmission electron microscopy.

They found that MoO_3 was first reduced to cubic oxygen-deficient Mo oxide (MoO_x) at a low temperature $(300^{\circ}C)$ due to the activation of H₂ by a trace amount of Rh. Then, <u>carbon atoms</u> could occupy oxygen vacancies to form cubic MoO_xC_y intermediates, which were essential to further transformation to α -MoC_{1-x} via the topological route. The final phase of carbides depended on the primary structure of initial intermediates and the route of structural evolution during the



carburization process, where a slow and stepwise reduction was necessary for the formation of active α -MoC_{1-x}.

"This work reveals the key factor of structural evolution during the carburization process," said Prof. Sun. "It provides a more facile and universal strategy for one-step synthesis of α -MoC_{1-x}."

More information: Xingtao Sun et al, In Situ Investigations on Structural Evolutions during the Facile Synthesis of Cubic α -MoC_{1-x} Catalysts, *Journal of the American Chemical Society* (2022). DOI: <u>10.1021/jacs.2c08979</u>

Provided by Chinese Academy of Sciences

Citation: Researchers reveal structural evolution during molybdenum carbides formation at atomic level (2022, December 1) retrieved 17 July 2024 from https://phys.org/news/2022-12-reveal-evolution-during molybdenum carbides formation at atomic level (2022, December 1) retrieved 17 July 2024 from https://phys.org/news/2022-12-reveal-evolution-during molybdenum-carbides-formation.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.