


Study reveals inconsistent effect of ramet connection in homogeneous environments

December 27 2022, by Zhang Nannan



The effect of stolon connection and water availability on the nine traits of *C. dactylon*. Credit: LU Zhijun

Physiological or clonal integration is a key ecological benefit where stolon or rhizome connections between ramets allow for translocation of resources from source-sites to sink-sites (in patchy or heterogeneous environments) within the clone. However, negative effects of ramet

connection have also been found in previous studies.

In order to address this controversial issue, researchers from the Wuhan Botanical Garden of the Chinese Academy of Sciences and University of Northern Colorado used *Cynodon dactylon* (Linn.) Pers. (Bermuda grass) as a [model plant](#) to determine whether ramet connection influences in homogeneous water availability (low/high), and whether the response could be attributed to the release of apical dominance or [root competition](#).

Results showed that stolon connection had no significant effects on the biomass of *C. dactylon*, while it affected plant morphology (stolon length) under the homogeneous high water condition. The apical dominance release treatment increased biomass and number of branches of *C. dactylon* distal sections demonstrating the effects of ramet separation. However, for apical sections, biomass, main stolon length, number of branches and inter- node length were significantly reduced following separation, resulting in the significant decrease of main stolon and inter- node length at the whole clonal fragment level.

The root competition experiment detected no significant differences of clonal fragments' performance between treatments (stolon connection and root competition), indicating weak root competition in homogeneous environments for *C. dactylon*, at least in early stages.

In conclusion, this study reveals that high water availability (favorable growth condition) and loss of ramet connection (acting as disturbance) would improve plant performance (escape strategy) via release of apical dominance or root competition (shaping plant performance) at early stages.

The paper is published in the journal *Flora*.

More information: Zhijun Lu et al, Revisiting ramet connection under homogeneous water availability environments, *Flora* (2022). [DOI: 10.1016/j.flora.2022.152163](https://doi.org/10.1016/j.flora.2022.152163)

Provided by Chinese Academy of Sciences

Citation: Study reveals inconsistent effect of ramet connection in homogeneous environments (2022, December 27) retrieved 3 May 2024 from <https://phys.org/news/2022-12-reveals-inconsistent-effect-ramet-homogeneous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.