

Why Asia-Pacific tropical seas are flush with marine life

January 20 2023

(a) Localized formation Sea surface Plankton bloom Converticine Converting Converting Converting REEF IS 20-40 km

Credit: Journal of Oceanography (2022). DOI: 10.1007/s10872-022-00673-2

A new Flinders University study has found why Southeast Asian seas produce more fish than the world's largest coastal upwelling regions combined.

While tropical oceans have been traditionally been considered low in both nutrients and phytoplankton productivity, Australian oceanographers have described this not the case, and how the effects of climate change may put this <u>natural process</u> at risk.



For the first time, they describe how transient short-lived wind events trigger the formation of significant phytoplankton blooms—the base of many marine food webs—near tropical reef islands, such as the disputed Spratly Islands in the South Island Sea and many other islands in the Pacific Ocean.

"The key here is that the tropical Pacific contains thousands of such islands, which implies that tropical wind events can create a large network of marine ecosystems existing year-round and spanning thousands of kilometers, possibly forming one of the largest upwelling biomes on Earth," says Associate Professor Jochen Kaempf, lead author of the new article published in the *Journal of Oceanography*.

Upwelling is a <u>physical process</u> that enriches surface layers of the ocean with dissolved nutrients where it fuels the production of phytoplankton.

Coastal winds are the principal driver of the most productive upwelling systems on Earth, such as those forming off the coasts of Peru-Chile and California, which are key global fish producers.

"We believe that this (Asia-Pacific) upwelling biome provides critical feeding habitats for migratory marine species such as <u>sea turtles</u>, tuna and whales.

"For instance, pygmy blue whales migrate from their feeding habitat in upwelling regions of southern Australia to their <u>breeding grounds</u> in Indonesian waters, where they seem to gather near Christmas Island and volcanic arc islands of the Banda Sea. Our findings suggest that this behavior is primarily feeding-related."

Associate Professor Kaempf says that the discovery paves the way for closer examination of phytoplankton bloom mechanisms in <u>tropical</u> <u>oceans</u>, particularly in the light of sea-level rise, enhanced wave erosion,



and changes in tropical rainfall and winds due to climate change.

The tropical Pacific Ocean contains most of the islands on Earth including hundreds of reef islands and atoll reefs.

More information: Jochen Kämpf et al, The island mass effect: a study of wind-driven nutrient upwelling around reef islands, *Journal of Oceanography* (2022). DOI: 10.1007/s10872-022-00673-2

Provided by Flinders University

Citation: Why Asia-Pacific tropical seas are flush with marine life (2023, January 20) retrieved 25 June 2024 from https://phys.org/news/2023-01-asia-pacific-tropical-seas-flush-marine.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.