


# Cells use mechanical principles to integrate within existing tissues, study shows

January 12 2023



The figure of the model shows that new cells integrate in vertices with more than three cells. Four or more cells (2) are more susceptible to opening up than points that connect three cells (1). Credit: Amin Doostmohammadi

Integration is not only a problem of social significance among humans, but also an issue for the cells that form us as humans. The addition of new cells into an already established group of cells is crucial for shaping the organs during normal development, but the very same mechanism is

also often hijacked by cancer cells as they spread within other cells.

A new interdisciplinary study reveals how new [cells](#) join a tissue by reading mechanical information from the neighboring tissue. They use cellular fingers called filopodia to touch the neighboring cells to open up the cell layer. The study is published in *Nature Communications*.

## **New cells use finger like structures to open up the layer of host cells**

To understand how new cells integrate within a cell layer, the group of scientists focused on the formation of cell layers in the . Frog cells share many characteristics with human cells, so they are useful for this type of research.

During embryo development, this multilayered cell assembly gets rearranged as new cells move from the to form tissue. This process must happen with accurate precision so that every single one of the incoming cells are properly placed.

High resolution microscope images showed that the incoming cells extend finger-like structures towards the cell layer that lies on top of them.

Detailed analyses of the experiments together with the theoretical model then revealed that the incoming cells use their finger-like extensions to pull on the vertices of the host layer, essentially probing for if they can open the vertex up and insert themselves.

Amin Doostmohammadi, assistant professor and the leader of the Active Intelligent Matter Group at the Niels Bohr Institute, explains how it works: "We found a particular protein complex that accumulates at the

vertex and allows the fingers from the incoming cells to make [physical contact](#) with the vertex. Its like the incoming cells are extending arms randomly to see where they can attach themselves."

## **Physical models predict the behavior of the living tissue**

"We then turned to a physical computer model of the cell layer to see if we could predict which vertices are more susceptible to these pulling forces," says Doostmohammadi. "That points that connect four or more cells are more susceptible to opening up than points that connect three cells."

"A similar trend was observed in the experiments, confirming the theoretical predictions, and showing that indeed incoming cells use pulling forces to find the weakest points within the host layer to insert themselves."

## **Physics and biological experiments go hand in hand**

The researchers used a very simple physics model of a network of vertices attached together through edges as a proxy for the complex network that cells form.

"With this simple model we could then test different types of vertices. We could pull on them to simulate the pulling forces that the new cells exert, we could check how much tension vertices can tolerate and whether they could indeed open up as a result."

"It was really a cross-talk between model and experiments, physics and biology: Designing a simple model, adjusting it to best represent key elements in the experiment, then making predictions that were verified

by the experiments."

## **Incoming cells create their own openings in existing cell layers**

However, at this point it was not clear how these weak points got created in the first place. "When there is no incoming cell, there are very few weak points in the cell layer and most of the vertices connect only three cells together. To our surprise we realized that exactly when incoming cells approach a cell layer a large number of vertices that connect four or five cells appear, suggesting that maybe the incoming cells themselves can affect the formation of weak points in the upper cell layer," says Doostmohammadi.

## **Mechanics guides cell integration**

These new findings highlight the importance of mechanical information in directing cell movement during [embryonic development](#) and provide a better understanding of how finger-like structures (filopodia), a common feature in invading cancers, are used to sense neighboring cells.

"It is amazing to see how mechanical this whole process is. Cells do not have a brain or a mechanism of decision-making. It is thus fascinating to find out how the incoming cells so accurately pull, probe, and choose the weak points in an existing cell layer, and if the environment is not favorable the cells even modify the layer and insert themselves," adds Doostmohammadi.

"The understanding gained from revealing the mechanisms of cell insertion has implications in situations where unwanted cells, such as [cancer cells](#), invade a cell layer. This understanding is a prerequisite for later development of therapies for treatment."

The work is published in the journal *Nature Communications*.

**More information:** Guilherme Ventura et al, Multiciliated cells use filopodia to probe tissue mechanics during epithelial integration in vivo, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-34165-0](https://doi.org/10.1038/s41467-022-34165-0)

Provided by University of Copenhagen

Citation: Cells use mechanical principles to integrate within existing tissues, study shows (2023, January 12) retrieved 29 June 2024 from <https://phys.org/news/2023-01-cells-mechanical-principles-tissues.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.