


Exotic wheat DNA helps breed 'climate-proof' crops

January 10 2023



Physiological assessment of HiBAP I panel, comparing elite and exotic-derived lines under heat stressed and yield potential conditions. **a** Number of lines from each group. **b** Effect of heat stress on yield (YLD), thousand grain weight (TGW), number of spikelets per spike (SPKLSP⁻¹), number of spikes per m² (SM2), plant height (Height), number of infertile spikelets per spike (infertile

SPKLSP⁻¹), harvest index (HI), grain weight per spike (GWSP), number of grains per spike (GSP), grain number (GM2), days to physiological maturity (DTM), days to anthesis or days to heading (DTA/DTH for yield potential and heat stress experiments respectively), and biomass at physiological maturity (BM_PM), showing the percentage difference compared to yield potential conditions. **c** Comparison of yield (YLD), thousand grain weight (TGW), grain number (GM2), biomass at physiological maturity (BM_PM), harvest index (HI), and Height between elite and exotic-derived lines in HiBAP I measured under both heat stress and yield potential conditions. The boxplots are defined as follows: center line = median; box limits = upper and lower quartiles, whiskers = 1.5x interquartile range; points = outliers. The significance of the difference between Elite ($n = 83$ biologically independent lines) and exotic-derived ($n = 66$ biologically independent lines) lines for each trait was assessed using two-tailed t tests with no assumption of equal variance. p -values below 0.01 were considered significant (*), below 0.001 very significant (**), and below 0.0001 highly significant (***). Credit: *Communications Biology* (2023). DOI: 10.1038/s42003-022-04325-5

Wheat containing exotic DNA from wild relatives benefits from up to 50 percent higher yields in hot weather compared with elite lines lacking these genes, according to a new study.

Following a year when temperature records have been smashed, research from the Earlham Institute in Norwich, in collaboration with the International Maize and Wheat Improvement Centre (CIMMYT), offers much-needed hope for improving crop resilience and [food security](#) in the face of climate change.

The field trials in Mexico also emphasize the importance of genetic diversity in key crops, where decades of selective breeding have reduced their ability to adapt to a rapidly warming planet.

There is growing uncertainty around the ability of major food crops to

continue to meet global demand as temperatures rise and weather events become more extreme.

Wheat provides more global calories than any other crop yet most of the wheat grown around the world has limited [genetic variation](#), making it vulnerable to the impacts of climate change.

"Wheat is responsible for around 20 percent of the calories consumed globally and is widely grown all over the world," says Professor Anthony Hall, study author and Group Leader at the Earlham Institute. "But we don't know whether the crops we're planting today will be able to cope with tomorrow's weather.

"To make matters worse, developing new varieties can take a decade or more so acting quickly is vital."

In collaboration with CIMMYT, Earlham Institute researchers set up a two-year field trial in Mexico's Sonora desert. They studied 149 wheat lines, ranging from widely-used elite lines to those selectively bred to include DNA from wild relatives and landraces from Mexico and India.

"Crossing elite lines with exotic material has its challenges," said Matthew Reynolds, co-author of the study and leader of Wheat Physiology at CIMMYT.

"There's a well-recognized risk of bringing in more undesirable than desirable traits, so this result represents a significant breakthrough in overcoming that barrier and the continued utilization of genetic resources to boost climate resilience."

The seeds were sown later in the season to force the plants to grow during hotter months, putting these crops under the kind of heat stress that is predicted to become the norm as global temperatures rise.

They found the plants bred with exotic DNA achieved a 50 percent higher yield over wheat without this DNA. Importantly, the exotic lines didn't perform any worse than the elite lines under normal conditions.

The researchers sequenced the plants to locate specific genetic differences responsible for the increased heat tolerance. They identified [genetic markers](#) that could allow the targeted introduction of this beneficial exotic DNA into elite lines, offering a quick way to improve climate resilience and mitigate against widespread crop failures.

Benedict Coombes, study author and Ph.D. student at the Earlham Institute, said, "As we try to produce more food from less land to feed a growing global population, we urgently need to future-proof the crops we're planting so they can thrive in an increasingly hostile climate."

"The key to this, we are increasingly finding, may lie within largely untapped genetic resources from wheat's wild relatives and landraces."

The researchers suggest breeding programs incorporate the heat tolerance traits as a pre-emptive strategy to produce wheat crops that can cope with a less predictable climate.

"This is science we can now use to make an impact almost immediately," added Professor Hall. "We've done the [field trials](#), we know what genetic markers we're looking for, and we're starting conversations with [wheat](#) breeders so this is hopefully going to be the first of many steps to contribute to global food security in the coming years."

"The discoveries we're making, and the action we're taking, will hopefully mean people around the world can continue to have nutritious food on their plates."

The work is published in the journal *Communications Biology*.

More information: Gemma Molero et al, Exotic alleles contribute to heat tolerance in wheat under field conditions, *Communications Biology* (2023). [DOI: 10.1038/s42003-022-04325-5](https://doi.org/10.1038/s42003-022-04325-5)

Provided by Earlham Institute

Citation: Exotic wheat DNA helps breed 'climate-proof' crops (2023, January 10) retrieved 29 April 2024 from <https://phys.org/news/2023-01-exotic-wheat-dna-climate-proof-crops.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.