

Study explores control options for black swallowwort

January 19 2023



Mean (+SE, back-transformed, n = 7) stem densities per square meter for Vincetoxicum nigrum >10-cm tall for seven treatments during a 3-yr study. Pretreatment values were in year 2018; values in 2019 and 2020 were 1-yr post first and second herbicide treatment, respectively. Bars within each year denoted by the same letter are not different (Fisher's protected LSD test with Bonferroni correction, P > 0.05). "ns" indicates no significant differences. Credit: *Invasive*



Plant Science and Management (2022). DOI: 10.1017/inp.2022.27

Black swallowwort is a difficult to control invasive vine that thrives in natural areas and perennial cropping systems across northeastern North America. To date, though, no scientific studies have been conducted to determine how the weed responds to common controls, such as mowing and broad-spectrum herbicides.

In an article featured in *Invasive Plant Science and Management*, researchers describe a three-year field study that begins to fill this information gap.

The team explored how black swallowwort responded to two glyphosate products and one triclopyr product. The weeds were either sprayed annually with a two percent solution of one the herbicides at flowering in early July—or were mowed in early July and then treated in late August.

The study showed the two glyphosate formulations were effective in reducing aboveground black swallowwort biomass, but they were less effective in reducing cover and stem densities. Researchers also determined that mowing failed to enhance the efficacy of the glyphosate applications on a consistent basis.

Triclopyr was generally ineffective in controlling black swallowwort—whether with or without mowing. The only significant effect of the <u>herbicide</u> was to increase grass cover in the test plots.

"Our study shows annual applications of glyphosate can be useful in managing infestations of black swallowwort vine," says Lindsey Milbrath of the USDA's Agricultural Research Service. "Further investigation will be needed to determine whether higher rates and more frequent



applications of triclopyr can also deliver effective control."

More information: Lindsey R. Milbrath et al, Response of black swallowwort (Vincetoxicum nigrum) to herbicides plus mowing, *Invasive Plant Science and Management* (2022). DOI: 10.1017/inp.2022.27

Provided by Cambridge University Press

Citation: Study explores control options for black swallowwort (2023, January 19) retrieved 6 May 2024 from <u>https://phys.org/news/2023-01-explores-options-black-swallowwort.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.