


Kenya's Rift Valley lakes are rising, putting thousands at risk. We now know why

January 17 2023, by Mathew Herrnegger



East African Rift Valley (left) and the study lakes in Kenya (right)

The East African Rift Valley sits between the Afar Triangle in Ethiopia and Mozambique. Within it lie a series of freshwater and alkaline lakes organized like a string of pearls.

The Kenyan lakes in the Rift have witnessed significant water level increases. Recent increases in lake areas range from 21% for Lake Naivasha to 123% for Solai. Homes, schools and hospitals have been flooded. A Kenyan government [report](#) estimates that about 80,000 households—400,000 people—have been affected by the floods since 2010. There are grave economic consequences, since agricultural fields and tourism infrastructure have been damaged.

To find out what is driving the rising [lake levels](#), my colleagues and I carried out a detailed analysis for several most affected lakes using satellite remote sensing data. We studied the changes of the lake surface areas, lake levels and water volumes, as well as the factors driving the inflows and the water balance of the lakes. These are chiefly [rainfall](#) and evaporation. [Our findings](#) provide compelling evidence that increases in rainfall since 2010 explain the lake level rises.

Human [activity](#) and changes in underground [permeability](#) have been advanced before to explain the lake level rises. The results of our study illustrate that surprisingly minor changes in the water balance are enough to explain the lake level rises. An increase of only 0.4–2% of mean annual effective rainfall leads to the observed phenomena.

Key findings on Rift lakes

Three key findings/trends emerged from our data analysis:

- Kenyan lakes showed low fluctuations and smaller water areas between 1984 and 2010, which have increased dramatically since 2010.

For instance, Lake Baringo had a [water area of 118km²](#) in 1995, the lowest value in 1984 to 2020. The water area expanded by over 52% in the years 2014 and 2020 to reach 195km².

The water area of Lake Solai, a smaller and less-known lake south of Lake Bogoria, increased from 3km² in 1984 to nearly 12km² in 2014 and 2020, a four-fold increase.

- There have been [dramatic fluctuations](#) between minimum and maximum water levels.

Significant differences between minimum and maximum water levels were found for all investigated lakes. The water level of Lake Solai increased by 8.5 meters, followed by Baringo (8.2 meters), Naivasha (7.5 meters), Nakuru (6.4 meters), Bogoria (5 meters) and Elementaita with 2.4 meters. For most lakes, the highest levels were found around 2013/2014—until 2020, in which new record maximums were observed.

To put this into perspective, the water level rise of over 8 meters between the 1980s and 1990s for Lake Baringo and Lake Solai is equivalent to the height of around two stories of a typical building.

- Rainfall has strongly increased in the lake catchments since 2010.

2009 was a breakpoint year for [lake catchment rainfall](#). The rainfall for the years 1981–2009 and 2010–2020 show noteworthy differences in the average annual rainfall values. This [matches well](#) with observations in lake levels, which showed low variability in water levels before 2009.

In the period 2010–2020, mean annual rainfall increased by up to 30% for the Baringo, Bogoria and Solai catchments. In Nakuru (21%), Elementaita (25%) and Naivasha (25%), the rainfall increases were somewhat lower. After 2018, annual rainfall increased by even more than 50% compared to the long term averages.

Another factor driving the water balance is evapotranspiration. This is

the amount of rainfall lost to the atmosphere. We found that this has also increased, though it's not clear by how much, since no data on the ground exists to validate the findings coming from numerical simulations.

Independent of the uncertainties regarding evaporation, our results suggest that increases in rainfall most likely explain the lake level rises.

Dealing with the consequences

The regional climate in Eastern Africa has shown significant variability in the past, with long lasting periods of severe drought but also wetter periods as we are currently observing. Water level fluctuations in the Rift Valley lakes are not new. [Lake Naivasha](#), for instance, was reduced to a puddle around 1850 and particularly low levels persisted throughout the 1940s and 1950s.

What is new is the immediate risk posed by rising lake levels. Compared to the past, much higher population densities around the lakes put more people, their homes and their crops in danger.

So will the lake levels keep rising? We don't know. Ten thousand years ago lake levels were significantly higher. South of Lake Turkana there was a lake called [Suguta](#) which was 300 meters deep, with a surface area of around 2,150km². This is over 10 times the size of current Lake Baringo. Lakes Bogoria and Baringo were connected and drained to the north towards Suguta, which abruptly disappeared around 8,500 years ago. (There are fears the lakes may [connect again](#), leading to a cross-contamination of alkaline and fresh water.)

There is little that humans can do to intervene, especially regarding increases in rainfall and change in climate. What we can do is to alleviate the impact and be prepared as a society. We can aim at holding back water in the catchments and upstream areas. Afforestation,

implementation of soil [water](#) conservation measures and enforcing infiltration of rainfall and surface runoff by technical and natural means are beneficial. Flood protection dams may be useful in some instances: this is however very costly.

Floods are natural phenomena, which were and will always be there. Preparedness is key in flood risk management. If Kenya and other countries affected by rising [lake](#) levels cannot change the situation, the only option is to learn to deal with it.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Kenya's Rift Valley lakes are rising, putting thousands at risk. We now know why (2023, January 17) retrieved 4 July 2024 from <https://phys.org/news/2023-01-kenya-rift-valley-lakes-thousands.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.