

NASA measures underground water flowing from Sierra to Central Valley

January 23 2023, by Jane J. Lee / Andrew Wang







River watersheds (outlined in various colors), mean rate of vertical motion from January 2006 to December 2021, and GPS sites (symbols). This study focuses on the Sacramento (blue), San Joaquin (green), and Tulare River (violet red) watersheds. The Klamath and north California Coast River watershed (pink) is also plotted. Color gradations depict nearly zero vertical motion (white color), slow uplift (light green), slow subsidence (light blue), and fast subsidence (magenta). See legend at bottom left for vertical rates. Contours of vertical motion at 10 mm/yr (violet to magenta) and 100 mm/yr in the southern Central Valley are labeled. The shape of the symbol at each GPS site designates the primary phenomenon deforming solid Earth at the site: porous response to groundwater change (red squares), elastic response to water oscillations (blue hexagons), and volcanic activity (green triangles). Early data before 2006 (gray circles) are omitted. The hexagons are filled with the color designating vertical motion in mm/yr observed at the GPS site (see legend). Credit: *Geophysical Research Letters* (2022). DOI: 10.1029/2022GL099583

In a recent study, scientists found that a previously unmeasured source—water percolating through soil and fractured rock below California's Sierra Nevada mountains—delivers an average of 4 million acre feet (5 cubic kilometers) of water to the state's Central Valley each year. This underground source accounts for about 10% of all the water that enters this highly productive farmland each year from every source (including river inflows and precipitation).

The Central Valley encompasses only 1% of U.S. farmland but produces 40% of the nation's table fruits, vegetables, and nuts annually. That's only possible because of intensive groundwater pumping for irrigation and river and stream flow captured in reservoirs. For at least 60 years, growers have been pumping more <u>water</u> from aquifers than can be



replenished by <u>natural sources</u>, causing the ground level to sink and requiring wells to be drilled deeper and deeper.

As water grows more scarce in the Central Valley due to <u>climate change</u> and <u>human use</u>, a more detailed understanding of the natural movement of groundwater offers a chance to better protect the remaining resources.

In the recently published study led by scientist Donald Argus of NASA's Jet Propulsion Laboratory in Southern California, researchers found that groundwater volume fluctuates more widely between dry and wet years than had been previously understood. In particular, the scientists observed a greater loss of groundwater during dry years than earlier studies had estimated. Argus and colleagues estimated that the Central Valley lost about 1.8 million acre feet (2.2 cubic kilometers) of groundwater per year between 2006 to 2021.

There is no way to directly measure the total volume of water on and under the Central Valley, but the satellites of the Gravity Recovery and Climate Experiment (GRACE) and GRACE Follow-On (GRACE FO) missions can accurately measure how much that volume changes from month to month. Argus and colleagues have been working for several years to combine such GRACE data with observations from a GPS research network that measures how land surfaces rise and subside. In central California, those motions are largely caused by increases and decreases in <u>underground water</u>.

Argus had previously used GPS to quantify the changing volume of water deep within the Sierra. For this new study, he and his co-authors used both GPS and GRACE measurements and subtracted the mountain groundwater change from the groundwater changes in both the mountains and the <u>valley</u> to obtain a more accurate estimate of change in the valley alone.



Then they compared that number with an estimate that co-author Sarfaraz Alam, a postdoctoral scholar at Stanford University, had calculated using a water-balance model. Such models attempt to account for all water entering and leaving an area from all processes, including river flows, precipitation, evaporation, and pumping from wells. The difference between the model's estimate and their new result, they believe, must come from the one process that was not included in the water-balance model: groundwater flowing from the mountains into the valley.

The volume of the inflow was surprising, according to Argus, because researchers believed they already had a good understanding of the amount of water entering and leaving the Central Valley. "We now know how much groundwater is going into and coming out of the aquifers during each season of the year, and during periods of drought and episodes of heavy precipitation," Argus said. This new understanding could be used, for example, to modify existing restrictions on watering during dry years versus wet years to better match usage with the available groundwater resources.

The paper is published in the journal Geophysical Research Letters.

More information: Donald F. Argus et al, Subsurface Water Flux in California's Central Valley and Its Source Watershed From Space Geodesy, *Geophysical Research Letters* (2022). DOI: 10.1029/2022GL099583

Provided by JPL/NASA

Citation: NASA measures underground water flowing from Sierra to Central Valley (2023, January 23) retrieved 26 April 2024 from <u>https://phys.org/news/2023-01-nasa-underground-</u>



sierra-central-valley.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.