


A new, better technology for X-ray laser pulses

January 18 2023



Edgar Kaksis (left) und Paolo Carpeggiani. Credit: Vienna University of Technology

The X-rays used to examine a broken leg in hospital are easy to produce. In industry, however, X-ray radiation of a completely different kind is

needed—namely, X-ray laser pulses that are as short and high-energy as possible. They are used, for example, in the production of nanostructures and electronic components, but also to monitor chemical reactions in real time.

Intense, extremely short-wave X-ray pulses in the nanometer [wavelength](#) range are difficult to produce, but now a new, simpler method has been developed at TU Wien (Vienna): the starting point is not a titanium-sapphire laser, which had mostly been used for this purpose, but an [ytterbium](#) laser. The crucial trick is that the light is then sent through a gas in order to change its properties.


Long wavelengths lead to short wavelengths

The wavelength of a laser beam depends on the material in which it is generated: In the atoms or molecules involved, electrons change from one state to another state with lower energy. This causes a photon to be emitted—its wavelength (and thus its color) depends on how much energy the electron lost during its change of state. In this way, different laser colors can be produced—from red to violet.

However, for creating laser beams with even much smaller wavelengths, special tricks have to be employed: First, [laser beams](#) with a long wavelength are created and shot at atoms. An electron is ripped away from the atom and accelerated in the electric field of the laser. It then turns back and collides again with the atom from which it came—and thereby it can produce short-wave X-rays. This technique is called "high harmonic generation."

"At first glance, the situation seems somewhat counter-intuitive," says Paolo Carpeggiani from the Institute of Photonics at TU Wien. "It turns out, in fact, that the larger the wavelength of the original laser beam, the smaller the wavelengths you can achieve in the end." However, the

efficiency of X-ray [radiation](#) production also decreases in the process: if you want to produce very short-wave radiation, its intensity becomes very low.



Graphical abstract. Credit: *ACS Photonics* (2022). DOI: 10.1021/acsp Photonics.2c01021

Ytterbium instead of titanium sapphire, gas instead of crystal

Until now, this technique was almost always employed by using titanium-sapphire lasers and then increasing the wavelength of their radiation with special crystals in order to generate the shortest possible X-ray radiation through High Harmonic Generation. However, the team at TU Wien has now developed a simpler and at the same time more powerful method: they used an ytterbium laser. Ytterbium lasers are simpler, cheaper and more powerful than titanium-sapphire lasers, but until now, their

performance in X-ray production was much lower.

At TU Wien, the wavelength of the ytterbium laser's radiation was first increased—not by sending this radiation through a crystal as usual, but by sending it through a molecular gas. "This increases the efficiency quite dramatically," says Paolo Carpeggiani. "Instead of the 20% we used to get, we get around 80%."

The paper is published in the journal *ACS Photonics*.

More information: Martin Dorner-Kirchner et al, HHG at the Carbon K-Edge Directly Driven by SRS Red-Shifted Pulses from an Ytterbium Amplifier, *ACS Photonics* (2022). [DOI: 10.1021/acsp Photonics.2c01021](https://doi.org/10.1021/acsp Photonics.2c01021)

Provided by Vienna University of Technology

Citation: A new, better technology for X-ray laser pulses (2023, January 18) retrieved 25 April 2024 from <https://phys.org/news/2023-01-technology-x-ray-laser-pulses.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.