

## **Review examines how to increase antidiabetic properties of purple vegetables and tubers**

February 15 2023



Graphical abstract. Credit: *Journal of Agricultural and Food Chemistry* (2022). DOI: 10.1021/acs.jafc.2c05879

The red, purple and blue pigments in fruits, vegetables and tubers called anthocyanins can reduce the risk of diabetes by affecting energy metabolism, gut microbiota, and inflammation. A new review article comparing the research results in the topic shows that the beneficial effect of anthocyanins on type 2 diabetes is increased if the anthocyanin is acylated, meaning that an acyl group is added to the sugar moieties of anthocyanin.



A great amount of acylated <u>anthocyanins</u> can be found in <u>purple potatoes</u>, , purple sweet potatoes, radishes, purple carrots and red cabbages, whereas bilberries and mulberries contain mostly nonacylated anthocyanins. Acylated anthocyanins are poorly absorbed in digestion, but they have probiotic properties and reduce the risk of diabetes more efficiently than nonacylated anthocyanins.

"The studies have shown that, in addition to changing physical and <u>chemical properties</u>, the acylation affects how the anthocyanins are absorbed and metabolized," says Postdoctoral Researcher Kang Chen at Food Sciences Unit, University of Turku, Finland.

The acylated anthocyanins are more effective antioxidants than the nonacylated anthocyanins, and they can also improve the intestinal barrier that enables the absorption of necessary nutrients. Furthermore, the acylated anthocyanins maintain <u>gut microbiota</u> homeostasis , suppress pro-inflammatory pathways, and modulate glucose and lipid metabolisms.

"The plant's genotype defines what kind of anthocyanins they produce. In general, purple vegetables contain many acylated anthocyanins. Also, purple potatoes, especially the Finnish variety called 'Synkeä Sakari,' is abundant in acylated anthocyanins," says Chen.

Acylated anthocyanins travel through our bodies from the upper gastrointestinal tract to the colon where they are metabolized by the gut microbiota. Glucose transporters are involved in anthocyanin absorption, but different <u>glucose</u> transporters are responsible for the absorption of acylated and nonacylated anthocyanins. The acylated and nonacylated anthocyanins also have different impacts on the enzymes involved in metabolism.

"The latest research has shown that the acylated and nonacylated



anthocyanins can impact type 2 diabetes in different ways," Chen summarizes.

The review article "Anthocyanins as Promising Molecules Affecting Energy Homeostasis, Inflammation, and Gut Microbiota in Type 2 Diabetes with Special Reference to Impact of Acylation" was published in the *Journal of Agricultural and Food Chemistry*.

**More information:** Kang Chen et al, Anthocyanins as Promising Molecules Affecting Energy Homeostasis, Inflammation, and Gut Microbiota in Type 2 Diabetes with Special Reference to Impact of Acylation, *Journal of Agricultural and Food Chemistry* (2022). DOI: 10.1021/acs.jafc.2c05879

Provided by University of Turku

Citation: Review examines how to increase antidiabetic properties of purple vegetables and tubers (2023, February 15) retrieved 20 June 2024 from <u>https://phys.org/news/2023-02-antidiabetic-properties-purple-vegetables-tubers.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.