

Coral reefs in the Eastern Pacific could survive into the 2060s, new study finds

February 13 2023



Ana Palacio, Ph.D., lead author of the study, surveys a coral reef in the eastern Pacific dominated by *Pocillopora* corals. Credit: Viktor Brandtneris

Scientists at the University of Miami Rosenstiel School of Marine, Atmospheric, and Earth Science found that some reefs in the tropical

Pacific Ocean could maintain high coral cover into the second half of this century by shuffling the symbiotic algae they host. The findings offer a ray of hope in an often-dire picture of the future of coral reefs worldwide.

While [global warming](#) is causing the loss of [coral reefs](#) globally, scientists believe that some corals are increasing their tolerance to heat by changing the symbiotic algae communities they host, which through photosynthesis provide them with the energy they need to live.

"Our results suggest that some reefs in the eastern tropical Pacific, which includes the Pacific coasts of Panama, Costa Rica, Mexico, and Colombia, might be able to maintain high coral cover through the 2060s," said coral biologist Ana Palacio-Castro, lead author of the study, alumna of the Rosenstiel School, and a postdoctoral associate at the school's Cooperative Institute for Marine and Atmospheric Studies. "However, while this may be seen as good news for these reefs, their survival may not continue past that date unless we reduce [global greenhouse gas emissions](#) and curtail global warming on a larger scale."

Shallow coral reefs in the eastern tropical Pacific Ocean are predominantly built by branching corals in the genus *Pocillopora*, which are extremely important for the reefs in the region. The microscopic algae they host in their tissue harvest light to help the coral produce energy to grow. The loss of these [symbiotic algae](#) causes the coral to turn white, or bleach, and the coral struggles to meet their energy needs, which can often prove fatal.

To better understand how corals improved their tolerance to heat stress, the researchers examined over 40 years' worth of coral reef-monitoring data from Panama, one of the longest datasets of its kind in the world. They analyzed temperature, coral cover, bleaching and mortality data spanning three ocean heatwaves—in 1982–1983, 1997–1998, and

2015–2016—along with data on algal symbiont community data during the last two.

The analysis showed that the 1982-83 heatwave significantly reduced coral cover on the reef, but the effects of the 1997-98 and 2015-16 El Niño were milder, especially for corals in the genus *Pocillopora*—sometimes known as cauliflower coral—the predominant reef-building coral in the eastern tropical Pacific. They also confirmed that during strong ocean heatwaves, the heat-tolerant alga *Durussdinium glynnii* becomes increasingly common in this particular lineage of corals, allowing them to better withstand periods of elevated temperatures. When combined with climate projections of future heat stress, the reefs that were predominantly composed of *Pocillopora* corals and that hosted this heat-tolerant alga were found to be better equipped to survive and maintain high levels of coral cover well into the second half of the current century, indicating that some [reef](#) systems may be more resilient to warming than previously thought.

"This study shows that there are some unusual reefs that may be able to survive for several decades as a result of their ability to shuffle symbionts," said Andrew Baker, professor of marine biology and ecology at the Rosenstiel School, and senior author of the study. "While we don't think that most reefs will be able to survive in this way, it does suggest that vestiges of our current reefs may persist for longer than we previously thought, although potentially with many fewer species. Coral reefs are incredibly valuable natural assets, providing coastal protection and fisheries benefits, and supporting many local communities. We can still make a difference by protecting them."

The study, titled "Increased dominance of heat-tolerant symbionts creates resilient [coral](#) reefs in near-term ocean warming," was published on Feb.13, 2023, in the journal *PNAS*.

More information: Palacio-Castro, Ana M., Increased dominance of heat-tolerant symbionts creates resilient coral reefs in near-term ocean warming, *Proceedings of the National Academy of Sciences* (2023). [DOI: 10.1073/pnas.2202388120](https://doi.org/10.1073/pnas.2202388120). doi.org/10.1073/pnas.2202388120

Provided by University of Miami Rosenstiel School of Marine, Atmospheric, and Earth Science

Citation: Coral reefs in the Eastern Pacific could survive into the 2060s, new study finds (2023, February 13) retrieved 26 June 2024 from <https://phys.org/news/2023-02-coral-reefs-eastern-pacific-survive.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--