

## Trained dogs can sniff out a deadly deer disease

February 8 2023, by Katherine Unger Baillie



Credit: Pixabay/CC0 Public Domain

Charlie, Jari, and Kiwi are pet dogs with a superpower: Their sensitive noses can distinguish between a healthy deer and one sick with chronic wasting disease (CWD), all from a whiff of the deer's poop.



That's the finding of a study by scientists from Penn's School of Veterinary Medicine, published in the journal *Prion*. Using feces samples from CWD-positive deer and deer in which CWD was not detected, the researchers trained the dogs to identify the odor of CWD, alerting their handlers to its presence in the lab and in the field.

"We were already quite certain that the dogs could detect the <u>volatile</u> <u>organic compounds</u> released by <u>chronic wasting disease</u> in feces," says Amritha Mallikarjun, a postdoctoral researcher in Penn Vet's Working Dog Center and lead author on the study. "Not only did we show this was possible, but we also answered a second, more interesting question, which is, Can they detect the disease in a simulated field setting, as they would if we were using the dogs to find the disease in the landscape of a forest or on a deer farm?"

The dogs indeed could, with enough accuracy to suggest that <u>detection</u> <u>dogs</u> could be a useful strategy in the fight to manage CWD.

"We learned a lot through the study and are now set up well to continue refining our training," says Cynthia Otto, the senior author on the study and director of the Working Dog Center.

CWD is a disease that affects a variety of deer species, including whitetailed deer, mule deer, and elk. Always fatal and highly contagious, the disease can hide away in an affected animal for a year or two before symptoms manifest: the deer losing weight—"wasting"—and developing neurological signs, like stumbling and drooling. No cure or treatment exists.

The disease has been in Pennsylvania since 2012, and the state has invested significantly in trying to contain it, with several tools in place for keeping tabs on its spread. One hurdle is figuring out which deer are affected. A healthy-looking but infected animal could shed prions,



malformed and infectious proteins, for many months or even years before succumbing to the illness. What's more, prions are known to bind to soil, potentially contaminating the land on which other animals may roam.

The gold-standard diagnostic test can only be performed after death by assessing an affected animal's brain. Some alternative tests have been trialed that involve taking a biopsy from a potentially infected animal while it's still alive, but deer are known to be highly stressed by being captured, and collecting these samples can be physically and logistically difficult for the people involved as well.

The Working Dog Center, the Pennsylvania Department of Agriculture, and the Wildlife Futures Program, a partnership between Penn Vet and the Pennsylvania Game Commission, were well positioned to try to contribute an additional technique for managing the disease: dogs and their highly sensitive noses. Ideally, dogs trained to discern CWDpositive from CWD-not-detected feces in a forest or deer farm could help state agencies and private landowners understand whether further testing or management would be needed to keep their land and herds free from the disease.

First, scientists had to prove the dogs could make this distinction reliably. The Working Dog Center began by enlisting three dogs from their citizen science program—Labrador retrievers Charlie and Kiwi and Finnish spitz Jari—to train on the Center's "scent wheel," a contraption equipped with eight ports, each containing a specific substance or scent.

The dogs proved adept at this task. Once they had been trained, using samples provided by the U.S. Department of Agriculture and processed by the Wildlife Futures Program, the dogs responded with great specificity, passing by the not-detected samples 90-95% of the time. Their sensitivity, however, was not as strong, alerting to just 40% of the



positive samples. Overall, the researchers found that the dogs spent more time at the ports containing positive samples than those with notdetected samples, suggesting that they perceived a difference even if they didn't always produce the trained response, such as a bark or sitting down when they smelled the positive sample.

Moving toward a more naturalistic setting, the researchers then experimented with having the dogs and their handlers try to discern CWD-positive samples placed throughout a large, privately owned field. To avoid contaminating the soil or having the dogs come in contact with the samples, two-gram samples of feces were placed in jars with mesh lids to allow the odor to waft out and then partially buried in the ground in different areas.

The researchers observed that the dogs responded to the positive samples more often than the not-detect samples in the field trial. In total, they detected eight of 11 positive samples in the field, with a false negative rate of 13%. Both handlers and dogs seemed to improve as they went, their accuracy increasing after their first field search.

"Given the amount of time that we trained these dogs and the novel environment, not to mention the fact that these are <u>pet dogs</u> and not trained search dogs, our results are promising," says Mallikarjun. "As we move forward and work with dogs that are specifically trained to search in a field setting and devote their entire lives to detecting this odor, they are going to do an even better job."

That's a step the Wildlife Futures Program is already taking, with canine handlers training "professional" detection dogs how to canvas fields and forests, searching for CWD.

The researchers believe that, while dogs don't represent a silver bullet in the fight against CWD, they may prove useful as an early warning



system, helping fill gaps in knowledge from other monitoring systems and management approaches.

"These dogs could increase the odds of catching an outbreak early," says Lisa Murphy, a study co-author and co-director of the Wildlife Futures Program.

The Penn Vet team is also collaborating with other groups not only to work with detection dogs but also to identify the odor dogs may be responding to in order to develop other systems for early detection. The lessons learned could be broadly useful.

"If we're able to tap into what we've learned with chronic wasting disease and apply it to other issues in agriculture and conservation, these <u>dogs</u> could be a major asset," Otto says.

**More information:** Amritha Mallikarjun et al, Canine detection of chronic wasting disease (CWD) in laboratory and field settings, *Prion* (2023). DOI: 10.1080/19336896.2023.2169519

## Provided by University of Pennsylvania

Citation: Trained dogs can sniff out a deadly deer disease (2023, February 8) retrieved 26 June 2024 from <u>https://phys.org/news/2023-02-dogs-deadly-deer-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.