


# London falcons ate fewer pigeons during lockdown

February 27 2023



London prey composition. Proportions of key prey items in London for peregrines during our study period. Starlings, pigeons and parakeets were the main prey groups; there was a lower proportion of pigeons consumed during lockdown, but starlings and parakeets were higher. Bar plots represent means of the respective prey groups from five sites, with error bars showing  $\pm 95%$  bootstrapped confidence intervals. Credit: *People and Nature* (2023). DOI: 10.1002/pan3.10445

Changes in peregrine falcon diets during COVID-19 lockdowns highlight the impact of human behavior on urban predators. The findings are from a new study published in the British Ecological Society journal, *People and Nature*.

In the study, researchers from King's College London and University of Bristol found that during lockdown, [peregrine falcons](#) in London were forced to change their diet away from [pigeons](#), with fewer of these birds being drawn in by human food supplies such as discarded [food waste](#) or direct feeding.

In the study, [citizen scientists](#) used online live streams to monitor 31 peregrine falcon nests in 27 U.K. cities over the course of three breeding seasons, the first of which took place during pandemic restrictions.

In London, peregrines took a lower proportion of pigeons as prey (-15%) and replaced them with starlings (+7%) and parakeets (+3%). However, in other cities, pigeons remained the dominant prey.

The study was written by Brandon Mak—a King's College London Ph.D. alongside Ed Drewitt—a University of Bristol ornithologist. Brandon Mak said, "Our results indicate that peregrines in larger, highly urbanized cities like London may be more dependent on, and hence more vulnerable to changes in, human activities which support their prey populations, particularly feral pigeons."

The changes to peregrine diets during the study raises questions surrounding how [pest control](#) may affect falcons and other predators that depend on "pest" species. For example, northern goshawk populations in Poland almost halved when farmers stopped rearing domestic pigeons and other poultry that would otherwise have been prey for them.

The management of pest species and their [food sources](#) are usually

human driven. Therefore reductions in pest species, like pigeons, can force raptors to switch prey or forage further away from their nests, which can result in poorer nutrition from less ideal prey, or a decrease in energy for fitness or reproduction due to the effort spent on hunting.

Brandon Mak explained, "The world is still learning about the consequences of lockdowns on wildlife, which promises to shed light on how human and animal lives are linked in our shared environments."

In the future, the authors of the study hope to contribute towards the Global Anthropause Raptor Research Network (GARRN) which brings together similarly conducted research from the pandemic.

"How did U.K. peregrines fare compared to elsewhere, and how did other raptors experience the lockdowns? We hope these questions will be answered in the coming years," added Mak.

**More information:** Brandon Mak et al, The raptor lockdown menu—Shifts in prey composition suggest urban peregrine diets are linked to human activities, *People and Nature* (2023). [DOI: 10.1002/pan3.10445](https://doi.org/10.1002/pan3.10445)

Provided by British Ecological Society

Citation: London falcons ate fewer pigeons during lockdown (2023, February 27) retrieved 20 April 2024 from <https://phys.org/news/2023-02-london-falcons-ate-pigeons-lockdown.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.