

Multifaceted analysis identifies transitional areas of vegetation as biodiversity hotspots

February 17 2023, by Zhang Nannan



Evergreen broad-leaved forest of Ailao Mountains in Yunnan. Credit: Li Jie

Climate change and biodiversity loss are interrelated environmental challenges facing the world today, and species living in transitional areas often show adaptive responses to climate change, making such areas



crucial for long-term biodiversity conservation. Therefore, it is essential to identify both transitional areas and biodiversity hotspots using multifaceted diversity measures to determine areas of overlap that require special conservation attention.

The cinnamon family, Lauraceae, is an important and diverse component of China's evergreen broad-leaved forests, making this family ideal for studying the relationships between areas of high biodiversity and transitional areas in order to identify multifaceted priority areas for conservation.

Accordingly, in a study published in *Ecological Indicators*, researchers from the Xishuangbanna Tropical Botanical Garden (XTBG) of the Chinese Academy of Sciences integrated <u>geographical distribution</u>, phylogenetic information, and functional diversity data to provide a systematic overview of Chinese Lauraceae.

The researchers first mapped and then identified spatial patterns of species richness, phylogenetic diversity, and functional diversity. Using these parameters, they then defined areas of neo- and paleo-endemism for Chinese Lauraceae and identified priority areas for conservation.

These multidimensional diversity patterns showed that for Chinese Lauraceae spatial patterns of species, evolutionary, and functional levels vary considerably among assemblages. Areas of paleo-endemism were identified in the border region of Yunnan and Guangxi, while neoendemic regions occurred mainly along the Yarlung Zangbo River and the Himalayas in southern Tibet.

Diversity hotspots were also found to overlap with evergreen broadleaved forests transition zones, highlighting the importance of transition zones for conservation in areas with diverse vegetation types.



"To increase conservation effectiveness under a changing climate, multidimensional assessments of diversity are desirable to improve the resolution of species patterns and provide better insights into both diversity and phylogenetically based conservation efforts," said Li Jie of XTBG.

More information: Run Zhou et al, Transitional areas of vegetation as biodiversity hotspots evidenced by multifaceted biodiversity analysis of a dominant group in Chinese evergreen broad-leaved forests, *Ecological Indicators* (2023). DOI: 10.1016/j.ecolind.2023.110001

Provided by Chinese Academy of Sciences

Citation: Multifaceted analysis identifies transitional areas of vegetation as biodiversity hotspots (2023, February 17) retrieved 23 June 2024 from <u>https://phys.org/news/2023-02-multifaceted-analysis-transitional-areas-vegetation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.