

NASA study focuses on impact effects on Mars rocks

February 16 2023, by Nick Oakes



This annotated image from NASA's Perseverance Mars rover shows its wheel tracks in Jezero Crater and a distant view of the first potential location it could deposit a group of sample tubes for possible future return to Earth. The image was taken on Aug. 29, 2022, the 542nd Martian day, or sol, of the rover's mission, by Perseverance's navigation camera. Sample tubes already filled with rock are currently stored in the rover's Sampling and Caching System. Perseverance will deposit select samples in designated locations. Subsequent NASA missions, in cooperation with ESA (European Space Agency), would collect these sealed samples from the surface and return them to Earth for indepth analysis as part of the Mars Sample Return campaign. Credit: NASA/JPL-Caltech



A NASA study describes how rocks could have been "shocked" and changed by meteorite impacts, once frequent on ancient Mars. This will improve our analysis of rock samples collected from the Red Planet.

"Because we're counting on these samples to reveal a record of Mars' geologic past, it would be important for us to understand if and how the rocks have been altered," said Dr. Svetlana Shkolyar, a <u>planetary</u> <u>scientist</u> at NASA's Goddard Space Flight Center in Greenbelt, Maryland. "The heat and pressure of an impact event can cause the rocks affected by it to melt. This means that when we study these rocks today, millions or even billions of years after the impacts occurred, the rocks could have had their original characteristics altered. It could even make one type of rock look like a different type of rock altogether."

Part of NASA's Perseverance rover's mission includes collecting <u>rock</u> <u>samples</u> from Mars that could later be retrieved and brought to Earth for study as part of the Mars Sample Return campaign. Scientists on the rover team want to make sure the most effective samples are chosen to accomplish two of the major goals of returned sample science—searching for signs of life and geologic history dating. Therefore, it is important for scientists to understand how these rock samples formed, and how they might have been altered over geologic time.





This image is of Jezero Crater on Mars, the landing site for NASA's Mars 2020 mission carrying the Perseverance rover. It was taken by instruments on NASA's Mars Reconnaissance Orbiter (MRO), which regularly takes images of potential landing sites for future missions. Jezero Crater sits within the Isidis Planitia region of Mars, where an ancient meteorite impact left behind a large crater some 750 miles (1,200 kilometers) across. This event is known as Isidis impact, and it forever changed the rock at the base of the crater. A later, smaller meteorite impact created the Jezero Crater within the Isidis impact basin. Scientists believe that these events likely created environments friendly to life. There is evidence of ancient river flow into Jezero, forming a delta that has long since been dry. Jezero Crater is thus likely to have been habitable in the distant past. Credit: NASA/JPL-Caltech/MSSS/JHU-APL



Shkolyar is lead author of a study published on May 3, 2022, in *Earth, Moon, and Planets* that describes how the rocks in these samples could be changed by impacts and how rover instruments can characterize those changes.

NASA, in partnership with ESA (the European Space Agency), plans to launch follow-up missions to the Mars 2020 Perseverance rover to retrieve the samples that have been selected and cached by Perseverance and bring them to Earth around 2033. This campaign is intended to allow far more <u>detailed studies</u> once samples are brought to Earth to gain a better understanding of Mars' past, present, and future, especially as it pertains to the possible presence of ancient life. Additionally, a key goal of the campaign is to help properly determine the age of events in Mars' geologic history. This is generally done by dating the rocks and using that to infer a chronology of Mars' history.

According to the study conducted by Shkolyar and her team, there are some important considerations to keep in mind when samples are selected. "For both astrobiology and dating the age of the rocks, the effect of impact shock is very important to consider," Shkolyar said. "When looking for fossil carbon in these rocks, which could be an indicator of past life, the alterations imposed by impact shock could alter the carbon signature." To ensure we understand the nature of the rocks, the Perseverance science team takes careful data to document the environmental context of samples before sampling and caching.





False-color topographic view of heavily cratered terrain on Mars from the Mars Orbiter Laser Altimeter instrument on board NASA's Mars Global Surveyor spacecraft. Red areas are high elevations and blue regions are low areas. The instrument collected altimetry data about the height of surface features on Mars until June 30, 2001. Credit: NASA/MOLA Science Team



The study focused on one kind of mineral, plagioclase feldspar, which is found in many rocks on Mars. The study details ways to recognize impact shock in plagioclase, if it occurred in the rocks studied, including improvement of the instrumental methods by which scientists characterize the chemistry of the rocks and identify evidence of shock in them. This way, when scientists cache and return those samples, they may find out whether the rocks have experienced impact shock, and how they might have been affected by this shock.

There is also much that can be learned from impact-shocked samples, such as adding precision to the chronology of the hypothesized Late Heavy Bombardment, a period early in the solar system's history when the planets may have been in less-stable orbits and could have been subjected to very frequent asteroid impacts.

"An example of how shocked samples could be beneficial would be to help us understand the history of impact on Mars," Shkolyar said. "Impact-shocked samples would contain geologic indicators which would help us to constrain the duration of the Heavy Bombardment more accurately."

More information: S. Shkolyar et al, Identifying Shocked Feldspar on Mars Using Perseverance Spectroscopic Instruments: Implications for Geochronology Studies on Returned Samples, *Earth, Moon, and Planets* (2022). DOI: 10.1007/s11038-022-09546-6

Provided by NASA's Goddard Space Flight Center

Citation: NASA study focuses on impact effects on Mars rocks (2023, February 16) retrieved 7 August 2024 from <u>https://phys.org/news/2023-02-nasa-focuses-impact-effects-mars.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.