

## How blocking cell wall formation stops bacterial cell division

March 23 2023



Bacterium Staphylococcus aureus under the microscope: Red fluorescent cell division protein FtsZ assembles so-called Z-ring in the center of the cell. Credit: University of Bonn/ Dominik Brajtenbach



Researchers still do not understand exactly how antibiotics kill bacteria. However, this understanding is necessary to develop new antibiotics. And that is precisely what is urgently needed, because bacteria are currently showing more and more resistance to existing antibiotics.

Researchers from the University Hospital Bonn (UKB) and the University of Bonn used high-performance microscopes to observe the effect of different antibiotics on the cell division of Staphylococcus aureus. They found that the biosynthesis of peptidoglycan, core component of the <u>bacterial cell wall</u>, is the driving force during the entire process of cell division. In addition, they clarified how exactly different antibiotics block cell division within a few minutes. The results have now been published in the journal *Science Advances*.

The bacterial cell wall maintains the shape and integrity of unicellular organisms. Cell wall synthesis plays a key role in bacterial growth: the cell division protein FtsZ forms the so-called Z-ring in the center of the cell, thus initiating the division process. A new cell wall is formed there, for which peptidoglycan is produced as the core component. This constriction thus gives rise to two identical daughter cells.

## Fluorescent proteins in Staphylococcus aureus under the microscope

The UKB research team led by Fabian Grein and Tanja Schneider, together with the team led by Ulrich Kubitscheck, Professor of Biophysical Chemistry at the University of Bonn, selected the bacterium Staphylococcus aureus, one of the most dangerous human pathogenic bacteria, as the model organism for their study. The focus was on the influence of antibiotics that inhibit peptidoglycan synthesis on cell division.



"We found a rapid and strong effect of oxacillin and the glycopeptide antibiotics vancomycin and telavacin on cell division. The cell division protein FtsZ served as a marker here and we monitored it," says Jan-Samuel Puls, a Ph.D. student at the Institute of Pharmaceutical Microbiology at UKB.

For this purpose, FtsZ was fluorescently labeled alongside other proteins. Then the researchers analyzed the effects on individual living bacterial cells over time and also used <u>super-resolution microscopy</u>. They established an automated image analysis for microscopy images that allowed them to quickly analyze all cells in the sample under study.

"Staphylococcus aureus is only about one micrometer, which is onethousandth of a millimeter. This makes microscopy particularly challenging," says Dr. Fabian Grein, junior research group leader at the UKB's Institute of Pharmaceutical Microbiology and a scientist at the German Center for Infection Research (DZIF).

## Antibiotic effect on cell wall biosynthesis machinery inhibits cell division immediately

The Bonn research team found that the formation of peptidoglycan is the driving force during the entire process of cell division. Previously, peptidoglycan synthesis was thought to be essential only during a specific part of this process. The team showed that inhibition of cell wall assembly by glycopeptide antibiotics in Staphylococcus aureus occurs rapidly and with a dramatic effect on cell division. In addition, they clarified in detail the specific role of essential penicillin-binding protein 2 (PBP2), which links cell wall components, in cell division. The  $\beta$ -lactam antibiotic oxacillin prevents the proper localization of this protein.



"This means that PBP2 does not get to the place where it is needed. As a result, the cell can't divide," Grein says. "Importantly, this all happens immediately after the antibiotics are added. So the first cellular effects, which have not been studied very intensively so far, are crucial." Therefore, in view of the alarming increase in <u>antibiotic resistance</u> worldwide, he hopes the study results will provide a better understanding of how exactly these agents work at the cellular level, and thus a key to the development of <u>new antibiotics</u>.

**More information:** Jan-Samuel Puls et al, Inhibition of peptidoglycan synthesis is sufficient for total arrest of staphylococcal cell division, *Science Advances* (2023). <u>DOI: 10.1126/sciadv.ade9023</u>

Provided by University of Bonn

Citation: How blocking cell wall formation stops bacterial cell division (2023, March 23) retrieved 13 May 2024 from <u>https://phys.org/news/2023-03-blocking-cell-wall-formation-bacterial.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.