

# Drones gather new and useful data for marine research, but they can disturb whales and dolphins

March 15 2023, by Jaclyn A. Aubin



Drones have allowed researchers to learn more about marine mammals. Credit: Jaclyn Aubin / GREMM, Author provided

Drones have changed the way researchers study whales and dolphins. While we were once confined to the decks of boats and observation platforms, glimpsing the backs of surfacing animals, we can now watch them from above. Gaining a bird's eye view of whales and dolphins has already taught us so much about their <u>physiology</u> and <u>behavior</u>.



However, there is a darker side to drone use in marine research.

My doctoral research investigates the behavior of beluga whales in the St. Lawrence Estuary. I have hundreds of hours under my belt as a drone pilot, flying over these amazing and endangered animals. As a Ph.D. student at the University of Windsor, I use drone footage of these whales to help us better understand their behavior and social structure, particularly differences between males and females.

#### **Observing behavior**

While using drones to conduct research, I quickly noticed that the drones we flew sometimes seemed to disturb the belugas. Other colleagues sometimes observed sudden dives, where most of the animals below the drone suddenly dove, often with considerable splashing. These responses appeared to be particularly common when the drone was flown at low altitudes, around 20 meters above the water.

We were concerned that this disturbance could impact our ability to study the whales, and worse, could have a negative impact on the whales themselves.

We conducted research into our observations, and our findings were <u>published in *Marine Mammal Science*</u>. We examined whether a series of variables relating to drone piloting impacted the likelihood of the belugas responding to the drone.

We predicted that responses to the drone would increase at low drone altitudes, when the drone's speed increased (increasing the noise of the rotors), when the drone approached the whales head-on, when <u>wind</u> <u>speeds</u> were low (making the drone more easy to hear), and on the first flight of the day.



We also examined variables relating to the whales being observed and predicted that drone disturbance would increase when the whales were in <u>small groups</u>, when offspring were present and when the whales were resting.

### **Drone altitudes**

We found that sudden dives were relatively rare, only occurring in about four percent of observations. However, sudden dives appeared to increase when drone altitude decreased. Sudden dives were particularly common when the drone was flown below 23 meters in altitude. This finding made sense to us: we had observed the drone was much more noticeable when flying directly above our heads, compared to when it was flying at 100 meters of altitude.

We also found that sudden dives appeared to be more common when flying over large groups. This finding was unexpected: earlier studies on dolphins have suggested that <u>small groups are more prone to disturbance</u>.

However, a <u>recently published study examining the impact of drones on</u> <u>bottlenose dolphins</u> found the same trend of disturbance increasing with group size. These authors suggested this is due to the "many eyes" effect, where large groups are more vigilant because there are more animals scanning for threats.

We also found that sudden dives commonly occurred when the drone first approached the whales. This suggests that the whales may be spooked by the initial approach of the drone.

## **Observer effects**

We also conducted a literature review examining the impact of drone



altitude on other whale and dolphin species. We found that drone disturbance rarely occurred when the drone was flown above 30 meters in altitude.

Interestingly, drone disturbance seemed more likely to be reported when authors included <u>detailed descriptions of how disturbance was assessed</u> and measured, suggesting that drone disturbance may be missed when researchers are not specifically looking for it.

We also found that most studies examining the impact of drones on whales and dolphins used small drones (less than five kilograms) which are much more discrete than the large, over 10 kilogram drone models that are frequently used in modern research programs.

#### **Future recommendations**

Our findings prompted us to make seven recommendations for future drone studies of whales and dolphins:

- 1. Drone studies of belugas should maintain an altitude of 25 meters or more.
- 2. Researchers using drones to study whales and dolphins should weigh the conservation benefit of low altitude flights against the potential for disturbance.
- 3. Pilots should use special caution when flying over large groups.
- 4. Pilots should use special caution when first approaching a group.
- 5. Pilots using large drones (those weighing more than 10 kilograms) should be particularly vigilant to drone disturbance



and should report on the effects of large <u>drones</u> on whales and dolphins.

- 6. Future drone studies should clearly define the disturbance behaviors they will watch for.
- 7. Using a precautionary approach, <u>drone</u> pilots should avoid sudden accelerations, avoid approaching animals head-on and maintain special caution in low wind conditions.

By reducing the disturbance impact of research on <u>whales</u> and dolphins, we can help protect these amazing animals.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Drones gather new and useful data for marine research, but they can disturb whales and dolphins (2023, March 15) retrieved 6 May 2024 from <u>https://phys.org/news/2023-03-drones-marine-disturb-whales-dolphins.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.