

## Drones detect moss beds and changes to Antarctica climate

March 6 2023



Credit: Queensland University of Technology

Researchers have been using drones to map large areas of Antarctica this summer in efforts to monitor the effects on vegetation due to climate change with the support of the Federal Government's Australian Antarctic Division.

The drone-derived imagery is being used by researchers from QUT and



Auckland University of Technology—with assistance from University of Wollongong—to evaluate the <u>fragile ecosystem</u>, particularly <u>moss</u> beds, and changes in the extreme environment.

For almost two months, the field team was based in Antarctica, with indications the <u>drones</u> they piloted captured unprecedented high-resolution imagery of vegetation and biodiversity in protected areas.

The vision taken from Australian Specially Protected Areas (ASPAs 135 and 136), not far from the scientists' base settlement of Casey Station, identified areas with moss and lichen not previously picked up by satellite.

QUT Center for Robotics researcher Dr. Juan Sandino, who specializes in mechatronics and automated remote sensing systems, helped develop and deploy the drones, classifying Antarctic vegetation at low altitude.

"Piloting these flights was at times challenging; however all the systems performed well under extreme cold conditions," he said.

He said working in Antarctica demanded rigorous physical preparation and had additional logistical pressure of ensuring the heavy equipment was operational.

The seven-year \$3.57 million project is co-led by QUT Professor Felipe Gonzalez and Auckland University of Technology Professor Barbara Bollard.

The project's key aims included monitoring the vegetation through smart sensors and <u>artificial intelligence</u>, modeling microclimates, and producing accurate maps of protected areas and other ice-free regions.

"Few <u>plants</u> can survive in Antarctica and mosses are the largest and



oldest plants to grow there with beds up to 50 meters wide and plants as old as 500 years," Professor Bollard said.

"The moss is sensitive to variations of temperature and moisture in Antarctica and is an important indicator of climate change."

Professor Gonzalez said mapping and monitoring the moss beds over time will help to understand the health of mosses and other vegetation.



Credit: Queensland University of Technology



Further drone-derived imagery as well as AI for other Antarctic regions may be used to evaluate the effectiveness of area-based conservation interventions in more parts of the icy continent.

QUT research, published recently in *Conservation Biology*, presented a case and a research agenda evaluating the conservation impact of Antarctica's protected areas.

Ph.D. researcher Joanna Burrows and supervisor Professor Kerrie Wilson, who are also part of SAEF, identified a gap in the research regarding the effectiveness of Antarctic Specially Protected Areas (ASPAs).

Burrows said that drones can provide new and more detailed information to evaluate the performance of ASPAs. "Drones could increase the likelihood of successful data collection by allowing remote and fragile ecosystems to be surveyed with low impact and can mitigate some accessibility issues," Burrows said.

**More information:** Joanna L. Burrows et al, Evaluating the conservation impact of Antarctica's protected areas, *Conservation Biology* (2023). DOI: 10.1111/cobi.14059

Provided by Queensland University of Technology

Citation: Drones detect moss beds and changes to Antarctica climate (2023, March 6) retrieved 26 April 2024 from <u>https://phys.org/news/2023-03-drones-moss-beds-antarctica-climate.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.