

# High winds can worsen pathogen spread at outdoor chicken farms

March 14 2023

---



Open environment chicken farms, which keep their flocks outdoors, often keep a variety of breeds, including heritage breeds. Credit: Olivia Smith, Washington State University

Farmers who keep their chickens outdoors may want to watch the weather. A study of chicken farms in the West found that high winds

increased the prevalence of *Campylobacter* in outdoor flocks, a bacterial pathogen in poultry that is the largest single cause of foodborne illness in the U.S.

Researchers found that about 26% of individual [chickens](#) had the pathogen at the "open environment" farms in the study, which included organic and free-range [chicken](#) farms. High winds the week prior to sampling and the farms' location in more intensive agricultural settings were linked to a greater prevalence of *Campylobacter*.

"Farmers need to be aware of the risk," said co-lead author Olivia Smith, a recent Washington State University Ph.D. graduate in the School of Biological Sciences. "These [environmental factors](#) are influencing if the poultry are going to have foodborne pathogens, so [farmers](#) need to be aware of what's around them. If there's a lot of wind and if they're in really agricultural areas, that's a problem."

To help reduce *Campylobacter* exposure, the researchers suggested farmers consider installing windbreaks and watch [weather patterns](#), so they can bring chickens inside during periods of [high winds](#) that could be blowing the bacteria onto their farms from nearby fields and livestock areas.

For the study, published in the [journal \*Animals\*](#), researchers tested chicken feces taken from 27 farms in California, Oregon, Washington and Idaho. They took samples at most of the farms once a year for three years. They found that the majority of the flocks at these farms, 69.4%, had some instance of campylobacter.





Open environment chicken farmers allow their flocks to roam outside as opposed to commercial producers who tend to keep poultry in fully enclosed barns. Credit: Olivia Smith, Washington State University

Researchers also interviewed the farmers about their management practices as well as types, breeds and ages of their chickens. Only 11 of the farms in the study were officially certified organic producers, but all of the farmers avoided using chemicals in their flocks, including antibiotics, vaccines or medications that kill parasites.

This is almost the exact opposite of commercial poultry producers who typically raise birds indoors in enclosed barns and treat them with medicines. These producers also select chicken breeds for efficiency,

such as "broiler" chickens that can grow fast and big providing a lot of meat, or "layer" breeds that can produce the most eggs, with the least amount of feed.

The growing market in local, organically minded food production has a different set of values, said Jeb Owen, a WSU entomologist and senior author on the paper. These farmers reduce or fully eliminate chemicals, keep a wide range of chicken breeds and allow their flocks to roam outside because they believe it is better for the animal and the environment. It is also what many consumers want, Owen said. But it doesn't come without risks.

"We've spent a century raising birds indoors and forgotten about all of these parasites and pathogens that chickens used to be afflicted with, but they didn't go away," he said. "Now you have this rapidly exploding market of producers who want to raise their birds outside, but they have no background knowledge of the disease risk."





A flock at an open environment chicken farm. Credit: Olivia Smith, Washington State University

Being outside means chickens are exposed to disease from [wild birds](#) and simply from contact with the ground, where they can pick up pathogens spread by feces of other infected birds.

Owen's lab has taken on a range of research to better understand the disease risk that faces open environment chicken farms, including a [study on enteric parasites](#) like worms that live in the birds' digestive systems and [another on ectoparasites](#), those that are found on the skin and feathers. His team is also undertaking a study to better understand the disease resilience of the many different breeds raised on these types of farms. The overall goal is to help farmers mitigate the risk.

"If they aren't doing it already, farmers should set up a professional relationship with a veterinarian to get their flocks checked and monitored on a regular basis," he said. "Whether for productivity or for animal welfare, you don't want your animals to be sick."

**More information:** Olivia M. Smith et al, Wind Speed and Landscape Context Mediate *Campylobacter* Risk among Poultry Reared in Open Environments, *Animals* (2023). [DOI: 10.3390/ani13030492](https://doi.org/10.3390/ani13030492)

Provided by Washington State University

Citation: High winds can worsen pathogen spread at outdoor chicken farms (2023, March 14) retrieved 16 June 2024 from <https://phys.org/news/2023-03-high-worsen-pathogen-outdoor-chicken.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--