

# How the 'marsupial sabertooth' *thylacosmilus* saw its world

March 21 2023

---



A reconstruction of *Thylacosmilus atrox*. Credit: © Jorge Blanco

A new study investigates how an extinct, carnivorous marsupial relative with canines so large they extended across the top of its skull could hunt effectively despite having wide-set eyes, like a cow or a horse. The

skulls of carnivores typically have forward-facing eye sockets, or orbits, which helps enable stereoscopic (3D) vision, a useful adaptation for judging the position of prey before pouncing. Scientists from the American Museum of Natural History and the Instituto Argentino de Nivología, Glaciología, y Ciencias Ambientales in Mendoza, Argentina, studied whether the "marsupial sabertooth" *Thylacosmilus atrox* could see in 3D at all. Their results are [published today](#) in the journal *Communications Biology*.

Popularly known as the "marsupial (or metatherian) sabertooth" because its extraordinarily large upper canines recall those of the more famous placental sabertooth that evolved in North America, *Thylacosmilus* lived in South America until its extinction about 3 million years ago. It was a member of Sparassodonta, a group of highly carnivorous mammals related to living marsupials. Although sparassodont species differed considerably in size—*Thylacosmilus* may have weighed as much as 100 kilograms (220 pounds)—the great majority resembled placental carnivores like cats and dogs in having forward-facing eyes and, presumably, full 3D vision. By contrast, the orbits of *Thylacosmilus*, a supposed hypercarnivore—an animal with a diet estimated to consist of at least 70 percent meat—were positioned like those of an ungulate, with orbits that face mostly laterally. In this situation, the visual fields do not overlap sufficiently for the brain to integrate them in 3D. Why would a hypercarnivore evolve such a peculiar adaptation? A team of researchers from Argentina and the United States set out to look for an explanation.

"You can't understand cranial organization in *Thylacosmilus* without first confronting those enormous canines," said lead author Charlène Gaillard, a Ph.D. student in the Instituto Argentino de Nivología, Glaciología, y Ciencias Ambientales (INAGLIA). "They weren't just large; they were ever-growing, to such an extent that the roots of the canines continued over the tops of their skulls. This had consequences, one of which was that no room was available for the orbits in the usual carnivore position

on the front of the face."

Gaillard used CT scanning and 3D virtual reconstructions to assess orbital organization in a number of fossil and modern mammals. She was able to determine how the visual system of *Thylacosmilus* would have compared to those in other carnivores or other mammals in general. Although low orbital convergence occurs in some modern carnivores, *Thylacosmilus* was extreme in this regard: it had an orbital convergence value as low as 35 degrees, compared to that of a typical predator, like a cat, at around 65 degrees.

However, good stereoscopic vision also relies on the degree of frontation, which is a measure of how the eyeballs are situated within the orbits. "*Thylacosmilus* was able to compensate for having its eyes on the side of its head by sticking its orbits out somewhat and orienting them almost vertically, to increase visual field overlap as much as possible," said co-author Analia M. Forasiepi, also in INAGLIA and a researcher in CONICET, the Argentinian science and research agency. "Even though its orbits were not favorably positioned for 3D vision, it could achieve about 70 percent of visual field overlap—evidently, enough to make it a successful active predator."



A reconstruction of the skull of *Thylacosmilus atrox*. Credit: © Jorge Blanco

"Compensation appears to be the key to understanding how the skull of *Thylacosmilus* was put together," said study co-author Ross D. E. MacPhee, a senior curator at the American Museum of Natural History. "In effect, the growth pattern of the canines during early cranial development would have displaced the orbits away from the front of the face, producing the result we see in adult skulls. The odd orientation of the orbits in *Thylacosmilus* actually represents a morphological compromise between the primary function of the cranium, which is to hold and protect the brain and sense organs, and a collateral function unique to this species, which was to provide enough room for the development of the enormous canines."

Lateral displacement of the orbits was not the only cranial modification that *Thylacosmilus* developed to accommodate its canines while retaining other functions. Placing the eyes on the side of the skull brings them close to the temporal chewing muscles, which might result in deformation during eating. To control for this, some mammals, including primates, have developed a bony structure that closes off the eye sockets from the side. *Thylacosmilus* did the same thing—another example of convergence among unrelated species.

This leaves a final question: What purpose would have been served by developing huge, ever-growing teeth that required re-engineering of the whole skull?

"It might have made predation easier in some unknown way," said Gaillard, "But, if so, why didn't any other sparassodont—or for that matter, any other mammalian carnivore—develop the same adaptation convergently? The canines of *Thylacosmilus* did not wear down, like the incisors of rodents. Instead, they just seem to have continued growing at the root, eventually extending almost to the rear of the skull."

Forasiepi underlined this point, saying, "To look for clear-cut adaptive explanations in evolutionary biology is fun but largely futile. One thing is clear: *Thylacosmilus* was not a freak of nature, but in its time and place it managed, apparently quite admirably, to survive as an ambush predator. We may view it as an anomaly because it doesn't fit within our preconceived categories of what a proper mammalian carnivore should look like, but evolution makes its own rules."

**More information:** Seeing through the eyes of the sabertooth *Thylacosmilus atrox* (Metatheria, Sparassodonta, *Communications Biology* (2023). [DOI: 10.1038/s42003-023-04624-5](https://doi.org/10.1038/s42003-023-04624-5)

Provided by American Museum of Natural History

Citation: How the 'marsupial sabertooth' thylacosmilus saw its world (2023, March 21) retrieved 13 March 2024 from

<https://phys.org/news/2023-03-marsupial-sabertooth-thylacosmilus-world.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--