

## Scientists explain how an infection can produce genetic diversity





Pathogen exposure suppresses avoidance of ascaroside pheromones. a, b, k, Schematics for bacteria exposure (a), quadrant assay on ascaroside pheromones (b), and histamine treatment during chemotaxis (k). c–j, Avoidance of the mixture of ascaroside pheromones ascr#2, ascr#3 and ascr#5 over a range of concentrations indicated for each ascaroside at equilibrium (c–g) or avoidance of individual ascarosides (h–j, 10 nM each at equilibrium) in wild-type adult hermaphrodites when exposed to E. coli OP50 or P. aeruginosa PA14. Positive avoidance index, avoidance. In box plots, the center line shows the median, box edges delineate 1st and 3rd quartiles and whiskers extend to minimum and maximum values. Numbers in parentheses, number of different assays. Dots, avoidance indexes of individual assays. P values are derived from two-tailed unpaired t test, asterisks indicate significant difference, \*\*\*\* P

## Citation: Scientists explain how an infection can produce genetic diversity (2023, March 6)



retrieved 25 June 2024 from <u>https://phys.org/news/2023-03-scientists-infection-genetic-diversity.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.