

# CryoSat reveals ice loss from glaciers

April 27 2023



While Greenland and Antarctica are losing masses of ice, so too are most of the glaciers around the world, but it's tricky to measure how much ice they are shedding. Thanks to ESA's CryoSat satellite and a breakthrough way of using its data, scientists have discovered that glaciers worldwide have shrunk by a total of 2% in just 10 years. That adds up to 2720 Gigatonnes in all. This can be imagined as a giant ice cube, bigger than Europe's highest mountain. Credit: ESA/Planetary Visions

When one thinks of the damage that climate change is doing, it's probable that what comes to mind is a vision of huge lumps of ice

dropping off one of the polar ice sheets and crashing into the ocean. While Greenland and Antarctica are losing masses of ice, so too are most of the glaciers around the world, but it's tricky to measure how much ice they are shedding.

Thanks to ESA's CryoSat satellite and a breakthrough way of using its data, scientists have discovered that glaciers worldwide have shrunk by a total of 2% in just 10 years, and it's because of higher air temperatures.

A paper, published in the journal *Geophysical Research Letters*, describes how scientists have used a particular technique of processing CryoSat data to reveal that glaciers lost a whopping 2,720 gigatons of ice between 2010 and 2020.

Their research also demonstrates that higher air temperatures are responsible for 89% of this ice loss.

Glaciers are found on all continents except Australia. They provide an essential source of freshwater. For example, glaciers in high-mountain Asia alone provide water for over 1.3 billion people. Glaciers are also important for industries such as hydropower.

The demise of glaciers around the world is, therefore, set to cause serious problems for local populations and those relying on outflow water further downstream.

In addition, ice being lost from glaciers is contributing more to [sea-level rise](#) than the ice being lost from either of the giant ice sheets on Greenland and Antarctica.

Despite mountain glaciers being key indicators of [climate change](#) and being important to society, estimates of global glacier mass loss have remained limited to a few scientific studies. This is because there are

numerous practical challenges in mapping and monitoring glaciers, which tend to lie in complex rugged terrain, and because there isn't a specific satellite mission dedicated to do this.

Noel Gourmelen, from the University of Edinburgh in the U.K., said, "I'm sure most people have seen photographs taken at different times that show how a glacier terminus has retreated over time. And we can see this from satellite images too.

"But we need to measure how a glacier's volume is changing to really make sense of what is happening."

The research team turned to ESA's CryoSat satellite, which carries a [radar altimeter](#) to measure the height of ice surfaces. This works well for measuring sea level and the height of sea ice, which is used to work out how ice thickness changes, and for measuring vast [polar ice sheets](#).

However, the footprint of this type of instrument is too coarse to measure and monitor mountain glaciers.

"Some years ago, we developed a technique of processing CryoSat data, called swath processing, which has revolutionized the use of CryoSat data over complex icy terrains. It unveils a wealth of new detail on glaciers," noted Dr. Gourmelen.

Livia Jakob, from Earthwave in the U.K., explained, "We've been able to use this technique to study glaciers all over the world and we can report that, in total, mountain glaciers lost 2% of their volume between 2010 and 2020.

"That adds up to 2,720 gigatons in all. This can be imagined as a giant ice cube, bigger than Europe's highest mountain, which is quite shocking. Importantly, we also found that air temperature, which causes

the ice surface to melt, accounts for 89% of this ice loss."

While warmer air temperatures are responsible for this decreasing "surface mass balance," the research team found that something called "ice discharge" was responsible for the other 11% of ice lost.

This is associated with glaciers that terminate at the coast, where warmer ocean waters are largely responsible for thinning the front of the ice flow.

Dr. Gourmelen said, "The relative contribution of decreasing surface mass balance and increasing ice discharge to sea-level change is well known for the Greenland and Antarctic ice sheets. Now we know more about how the atmosphere and ocean are teaming up to melt glaciers. There is still plenty of work to do to refine these numbers, and to incorporate this knowledge into our glacier projections."

Stephen Plummer from ESA explains, "The innovative work using CryoSat in swath altimetry mode demonstrates the value of its altimeter for monitoring glaciers, thus achieving one of the mission's secondary objectives.

"This work laid the basis for the Glacier Mass Balance intercomparison Exercise, Glambie, to reconcile the different estimates of glacier mass balance from a multitude of satellite and in-situ methods. It also helps in guiding the design of the Copernicus Sentinel Expansion CRISTAL mission for monitoring land ice, ensuring continuity in monitoring [glaciers](#) globally."

**More information:** Livia Jakob et al, Glacier Mass Loss Between 2010 and 2020 Dominated by Atmospheric Forcing, *Geophysical Research Letters* (2023). [DOI: 10.1029/2023GL102954](https://doi.org/10.1029/2023GL102954)

Provided by European Space Agency

Citation: CryoSat reveals ice loss from glaciers (2023, April 27) retrieved 26 April 2024 from <https://phys.org/news/2023-04-cryosat-reveals-ice-loss-glaciers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.