

## **Evolution of two contagious cancers affecting Tasmanian devils underlines unpredictability of disease threat**

April 20 2023



Tasmanian devil (Freycinet, Tasmania). Credit: Max Stammnitz

Transmissible cancers, which occur only rarely in the animal kingdom, are spread by the transfer of living cancer cells. In the case of Tasmanian



devils, the cells are transferred through biting—a behavior that is common in devils especially in fights over mates and food.

Tasmanian devils are susceptible to two fatal transmissible cancers called devil facial tumor 1 (DFT1) and devil facial tumor 2 (DFT2) that have caused rapid population decline in recent decades. The two cancers both manifest with disfiguring facial tumors.

In a new study, University of Cambridge researchers mapped the emergence and <u>mutations</u> of DFT1 and DFT2 and characterized these cancers' ongoing evolution. The findings underline the continued threat that transmissible cancers pose to Tasmanian devils. The results are published today (April 20) in the journal *Science*.

"The incredible fact that Tasmanian devils have not one, but two, transmissible cancers, makes it possible to compare their evolution, and this gives us new insights into the key mechanisms involved," said lead author Elizabeth Murchison, Professor of Comparative Oncology and Genetics at the Department of Veterinary Medicine, University of Cambridge.

"By looking at the mutations that have accumulated in these cancers' DNA, we can trace the origins and evolution of these diseases. Our results show that the two cancers arose through similar processes and that both have striking signals of ongoing evolution. It is difficult to predict how this continued cancer evolution will impact devils."





Tasmanian devil with a facial tumor (Freycinet, Tasmania). Credit: Max Stammnitz

The researchers created an improved "reference genome"—essentially a map of the entire DNA sequence—of the Tasmanian devil and compared this to DNA taken from 119 DFT1 and DFT2 tumors. DFT1 was first observed in 1996 in Tasmania's northeast and is now widespread throughout Tasmania. DFT2, on the other hand, was first observed in 2014 and remains confined to a small area in Tasmania's southeast. The scientists identified mutations in the tumors and used these to build "family trees" of how the two cancers had each independently arisen and evolved over time.

By tracking mutations the researchers discovered that DFT2 acquired



mutations about three times faster than DFT1. As mutations usually occur during <u>cell division</u>, the most likely explanation is that DFT2 is a faster growing cancer than DFT1, say the researchers, underlining the importance of DFT2 as a threat.

"DFT2 is still not widespread in the devil population, and very little is known about it. We were really startled to see just how quickly it was mutating, alerting us to what could be a very unpredictable threat to the devils in the long term," said Maximilian Stammitz, first author of the study.

The team found that DFT1 arose in the 1980s, up to 14 years before it was first observed, whereas DFT2 emerged between 2009 and 2012, only shortly before it was detected.

Mapping the mutations revealed that DFT1 underwent an explosive transmission event shortly after it emerged. This involved a single infected devil transmitting its tumor to at least six recipient devils.





Tasmanian devil (Freycinet, Tasmania). Credit: Max Stammnitz

DFT1 has now spread throughout almost the entire devil population and has <u>recently been reported in the far northwest of Tasmania</u>, one of the few remaining disease-free regions of the state.

Researchers also identified for the first time an instance of DFT1 transmission between a mother and the young in her pouch. Additionally, they found that the <u>incubation period</u>—the time between infection and the emergence of symptoms—can in some cases be a year or more. These findings have important implications for conservation scientists working to protect the species.



"I come from Tasmania and love Tasmanian devils—they have a special place in my heart," said Murchison. "Transmissible cancers pose an unprecedented and unpredictable threat to Tasmanian devils. This research highlights the continuing importance of monitoring and conservation programs. It also gives us new insights into the evolutionary mechanisms operating in cancer more broadly, including in human cancers."

**More information:** Maximilian R. Stammnitz et al, The evolution of two transmissible cancers in Tasmanian devils, *Science* (2023). DOI: 10.1126/science.abq6453. www.science.org/doi/10.1126/science.abq6453

## Provided by University of Cambridge

Citation: Evolution of two contagious cancers affecting Tasmanian devils underlines unpredictability of disease threat (2023, April 20) retrieved 23 June 2024 from <u>https://phys.org/news/2023-04-evolution-contagious-cancers-affecting-tasmanian.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.