

## **Researchers image magnetic behavior at the smallest scales to date**

April 12 2023, by Michael O'Boyle



4D-STEM performed on a sample of iron arsenide. An electron beam from the top is directed into a sample of the antiferromagnet. The effects of the magnetic order on the beam pattern are observed in the detector at the bottom. Credit: The Grainger College of Engineering at the University of Illinois Urbana-Champaign



Permanent magnets, the kind found on refrigerators everywhere, exist because their constituent atoms behave as miniature magnets. They align and combine to form the larger magnet in a phenomenon called ferromagnetism. There are some materials where the atomic magnets instead form an alternating pattern, so the material has no net magnetization. Such antiferromagnets have attracted attention for their potential to create faster and more compact magnetic memory devices for computing.

Realizing the full potential of antiferromagnetic devices will require sensing their atom-to-atom magnetic patterns, something that has not yet been achieved. However, researchers at the University of Illinois Urbana-Champaign, led by Pinshane Huang, a professor of materials science & engineering, have made progress toward this goal. In the journal *Ultramicroscopy*, they report a new electron microscopy technique that can resolve magnetic behavior on the scale of angstroms—tenths of nanometers, nearly on the scale of individual atoms. They use this technique to fully resolve the antiferromagnetic order in iron arsenide for the first time.

"We are working to develop new techniques that can resolve <u>magnetic</u> <u>behavior</u> of individual atoms, and this study is an important step," Huang said. "The best techniques before now have achieved resolutions of a few nanometers. We have vastly exceeded that record."

Microscopic magnetism is often measured with scanning <u>transmission</u> <u>electron microscopy</u>, or STEM, in which an <u>electron beam</u> is focused into a material. The electric interactions between the beam and the material's structure have been famously used to provide images of individual atoms in the material, but the beam also interacts with the material's magnetic structure.

Although this much weaker interaction is enough to determine the longer-



range magnetic order found in ferromagnets, a far more precise technique is needed to observe the atom-to-atom variation in antiferromagnets.

"In standard, low-resolution STEM experiments, the magnetic interaction can be understood as a small deflection of the electron beam," said Kayla Nguyen, a postdoctoral research associate and co-lead author of the study. "On the scale of individual atoms, though, this picture breaks down. The magnetic interactions create complicated and subtle signals in the beam pattern which require new tools to analyze and understand."

To achieve higher resolution, the researchers turned to a more powerful method called four-dimensional STEM. Standard STEM techniques record drops in the beam's intensity as it interacts with the material, but 4D-STEM captures full two-dimensional scattering patterns as the electron beam scans along the two directions of the material's surface (for four-dimensional data). These data allowed the researchers to search the full beam patterns for the more intricate signals of atomic antiferromagnetism.

A crucial step of the analysis was simulating the magnetic fields within the iron arsenide sample, for which the researchers wrote a <u>software</u> <u>package</u> called Magnstem. Graduate student and co-lead author Jeffrey Huang explained that the package allowed them to add <u>magnetic effects</u> specific to their material and study the effects they had on electron beam patterns.

"Magnstem simulations allowed us to compare the electron patterns with magnetic effects turned on versus turned off, something that would be quite difficult to do in a real experiment," he said. "We saw that the effects of the magnetic and electric signals occur on different parts of the pattern and can be extracted separately."



By combining 4D-STEM with Magnstem simulations, the researchers resolved magnetic order down to 6 angstroms. While this does not resolve magnetic effects on the scale of individual atoms, it allowed them to resolve the antiferromagnetic pattern of iron arsenide, which repeats in cells of 12 atoms.

"Our work has shown it is possible to resolve small-scale magnetic order in electron microscopy experiments and in simulations, almost at atomic resolution," Pinshane Huang said. "We are actively developing techniques that will build on this result."

The researchers worked in collaboration with the research groups of Daniel Shoemaker and André Schleife, both professors of materials science & engineering. Shoemaker's group produced samples of iron arsenide, and Schleife's group performed simulations of the material's magnetic structure.

**More information:** Kayla X. Nguyen et al, Angstrom-scale imaging of magnetization in antiferromagnetic Fe2As via 4D-STEM, *Ultramicroscopy* (2023). DOI: 10.1016/j.ultramic.2023.113696

Provided by University of Illinois Grainger College of Engineering

Citation: Researchers image magnetic behavior at the smallest scales to date (2023, April 12) retrieved 28 June 2024 from <u>https://phys.org/news/2023-04-image-magnetic-behavior-smallest-scales.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.