


Multicellular organisms without a nervous system behave like a swarm, finds study

April 20 2023



Partial cross-section of *T. adhaerens*. The animal consists of three layers of cells: a dorsal and ventral epithelium layer, with an inner layer of fiber cells. The dorsal and ventral cells are ciliated, with the ventral cilia providing propulsive force for crawling along surfaces, the animal's primary form of locomotion. Not shown are other specialized cells, such as digestive and crystal cells, that are not directly responsible for movement. Credit: *Proceedings of the National Academy of Sciences* (2023). DOI: 10.1073/pnas.2206163120

Coordination between individuals is a fundamental challenge in collectives, and this is just as true for a country's government as it is for simple multicellular organisms. How does a group of many cells coordinate itself into a body that performs coherent behavior? In more

evolved animal groups, that's the nervous system's job.

But if we go back to the beginning of the evolutionary tree and take a look at the first multicellular animals, we'll notice that some of them are capable of moving in a coordinated manner even though they don't have nervous systems. One example is the placozoan *Trichoplax adhaerens*, the simplest multicellular animal organism ever described. *T. adhaerens* is a type of marine amoeba with a body so undifferentiated that it not only lacks a nervous system, but also a mouth, stomach, muscles, blood, or veins. Yet, it is motile thanks to the [collective action](#) of thousands of cilia, small, hairlike structures that can beat in any direction, allowing motion.

This caused researchers to wonder: how can a collection of cells move in a seemingly coordinated way if it doesn't have a "brain?" In a new paper published in *Proceedings of the National Academy of Sciences*, Science of Intelligence PI Pawel Romanczuk, together with Mircea Davidescu (Princeton University), Thomas Gregor (Princeton University), and Iain Couzin (University of Konstanz), investigated coordination among cells in *T. adhaerens* by measuring how well the collective moves together in [different individuals](#) of varying sizes.

They found that the cells that compose the collective will initiate motion when the cells find themselves right at the border between a state of order and a state of chaos—that is, when they are in the so-called "state of criticality." This phenomenon, which is known to occur in the animal brain, has also been observed in a Mexican fish collective described in [another paper](#), published in *Nature Physics*. In that study, the researchers had observed the fish school perform fascinated, wave-like motions to escape predators whenever the group reached the stage of criticality, and the researchers had described that as a sort of "collective brain."

The current study combined detailed observation of *T. adhaerens* with a

[mathematical model](#) that simulated its movement in the computer. Tracking under the microscope allowed to quantify how different parts of individual animals move with respect to each other when they explore their environment. Mathematically speaking, the simple multicellular animals were modeled as a self-organized collective of motile cells, i.e., cells able to move, coupled to each other. In [computer simulations](#), researchers were able to generate predictions for the observed behaviors, and found that the behavior of the animals can be only explained if they operate at criticality.

If criticality is what allows *Trichoplax adherens* to coordinate the motion of its different parts, it also comes at a cost, and therefore the larger the animal (seen as a collective of cells,) the less organized its locomotion will appear. The scientists also went a step further and quantified the trade-off between increased size and motion coordination in *T. adherens*, and argued that there is a maximum size (1–2 mm) beyond which it is no longer possible for the system to move in a coordinated manner. This could shed new light on how and why larger organisms evolved hierarchical structures, such as nervous systems.

More information: Mircea R. Davidescu et al, Growth produces coordination trade-offs in *Trichoplax adhaerens* , an animal lacking a central nervous system, *Proceedings of the National Academy of Sciences* (2023). [DOI: 10.1073/pnas.2206163120](https://doi.org/10.1073/pnas.2206163120)

Provided by Technische Universität Berlin

Citation: Multicellular organisms without a nervous system behave like a swarm, finds study (2023, April 20) retrieved 3 May 2024 from <https://phys.org/news/2023-04-multicellular-nervous-swarm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.