


Novel nanocages for delivery of small interfering RNAs

April 19 2023



Schematic representation of the use of nanocages for siRNA delivery. When binding siRNA (arrow A), the nanocages with a typical diameter of 5-6 nanometre cluster into cage-RNA nanoparticles with a diameter of some 150 nanometres. This is small enough to enter a cell (arrow B) and deliver the siRNA. Image taken from the paper. Credit: *Chem* (2023). DOI: 10.1016/j.chempr.2023.03.018

Small interfering RNAs (siRNAs) are novel therapeutics that can be used to treat a wide range of diseases. This has led to a growing demand for selective, efficient, and safe ways of delivering siRNA in cells. Now, in a cooperation between the Universities of Amsterdam and Leiden, researchers have developed dedicated molecular nanocages for siRNA delivery. In a paper just out in the journal *Chem* they present nanocages


that are easy to prepare and display tunable siRNA delivery characteristics.

The nanocages were developed in the research group for Homogeneous, Supramolecular and Bio-inspired catalysis of Prof. Joost Reek and Bas de Bruin at the University of Amsterdam's Van 't Hoff Institute for Molecular Sciences, and further studies in the group Prof. Alexander Kros at the Leiden Institute of Chemistry.

The researchers were motivated by the potential of siRNA in [gene therapy](#), which requires the need for effective delivery systems. They set out to develop nanocages with [functional groups](#) at the outside, making the cages capable of binding siRNA strands. As the binding is based on reversible bonds, the siRNA can in principle be released in a cellular environment. To explore the delivery characteristics of their nanocages, the researchers performed a laboratory study using various human cancer cells.

A range of nanocages

The nanocages are constructions of small molecular building blocks, so-called ditopic ligands, that are connected using metal atoms. A typical cage consists of 12 metal atoms and 24 ligands, hence the abbreviation M12L24. The researchers designed and synthesized five different ligands to form molecular cages with different siRNA binding affinities. They then prepared a range of siRNA binding nanocages using platinum or palladium as connecting metal. The palladium nanocages are less stable in a cellular environment, and decomposition is one of the siRNA releasing mechanisms.



Four Pt₁₂L₂₄ nanocage structures investigated in the study, using platinum and a variety of ligands. The first nanocage was also synthesised in a variant using palladium instead of platinum. Image taken from the paper. Credit: University of Amsterdam

After screening [nanocage](#) characteristics such as stability and siRNA binding capability, the delivery characteristics were put to the test in assays based on siRNA-mediated Green Fluorescent Protein (GFP) silencing. The cages were used to deliver siRNA to human GFP-expressing cells, so that fluorescence measurements could establish successful siRNA delivery. Two types of human cell lines were used: HeLa and U2Os.

Cage composition determines siRNA delivery

To their surprise, the researchers could not only demonstrate satisfying siRNA delivery, but also discovered a remarkable differentiation depending on the metal used in the nanocage. Where a platinum-based Pt₁₂L₂₄ nanocage showed highly effective siRNA delivery to U2OS cells, it showed little efficiency for HeLa. By contrast, the palladium-based Pd₁₂L₂₄ nanocage, derived from the same ligand building block,

delivered siRNA to HeLa but not to U2OS. Such differentiation could not be observed in experiments where a commercially applied delivery system (lipofectamine) was used. The M12L24 nanocages thus introduce the possibility of tuning siRNA delivery characteristics by tuning the nanocage composition.

In their *Chem* paper, the researchers consider this unique cell selectivity feature of the nanoparticles a promising addition to the field of targeted RNA gene material delivery, whose full potential is yet to be uncovered. Even though the current results were obtained in highly controlled laboratory research, they expect that the tunable RNA delivery of their nanocages will spawn future developments of highly desirable selective RNA nanomedicines.

More information: Eduard O. Bobylev et al, The application of M12L24 nanocages as cell-specific siRNA delivery agents in vitro, *Chem* (2023). [DOI: 10.1016/j.chempr.2023.03.018](https://doi.org/10.1016/j.chempr.2023.03.018)

Provided by University of Amsterdam

Citation: Novel nanocages for delivery of small interfering RNAs (2023, April 19) retrieved 11 July 2024 from <https://phys.org/news/2023-04-nanocages-delivery-small-interfering-rnas.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--