


Better recycling of plastic packaging: New process extracts fragrances

April 20 2023



Principle of the multivariate analysis of a set of FTIR spectra. The color scale corresponds both to the intensity and the percentage of HDPE and the extractables (odorants, impurities). Credit: Fraunhofer LBF

What doesn't smell good is hard to recycle. This simple rule also applies to the growing volume of plastic waste worldwide. One way to recycle it in an environmentally compatible and climate-friendly manner as high-quality post-consumer recyclates is through improved sorting and

reprocessing. Until now, the reduced material quality has considerably limited the reuse of plastic recyclates, and this is mainly due to their odor.

Scientists at the Fraunhofer Institute for Structural Durability and System Reliability LBF have now developed a new environmentally friendly process on a laboratory scale to remove odors from [plastic](#) packaging.

The new process is based on pressurized water [extraction](#). It removes the tracer fragrance limonene from commercial HDPE packaging and does not require organic solvents. This reduces costs and protects the environment. In this way, the material quality of processed plastic waste can be significantly increased within one hour.

The in-process analysis is based on [infrared spectroscopy](#) and [mass spectrometry](#) that provide the research team with data on the chemical composition of the samples as a function of different extraction conditions. The analytical data show that significantly less limonene is present in the samples after extraction. Furthermore, in addition to the fragrance, other impurities and short-chain HDPE are removed from the samples that were originally contained in the packaging.

Based on these analyses, the Darmstadt experts determined the optimal process parameters for pressurized water extraction of fragrances from HDPE packaging. "The project results demonstrate the benefits of a systemic approach to solving current plastics technology issues with great social relevance," emphasizes Dr. Guru Geertz, who oversees the project at Fraunhofer LBF.

Material analysis with machine learning methods optimizes extraction process

To develop the process, detailed insights into the chemical kinetics of the extraction process were necessary, which were made possible by a novel approach for in-process analytics: Evaluating the data with the aid of machine learning methods was the key to optimize the extraction parameters in terms of the desired economical process control. At the current stage of development, an application scenario for the new process is emerging for the improved treatment of plastic waste.

"The extraction process we have developed shows a way to reprocessed single-use plastics with an increased range of applications, and this serves to protect the environment," says Dr. Geertz. Due to the underlying concept, the process is equally suitable for industrialized, emerging, and developing countries, so that all market participants who bring plastic products into the trade can potentially benefit from it.

Provided by Fraunhofer USA

Citation: Better recycling of plastic packaging: New process extracts fragrances (2023, April 20) retrieved 10 July 2024 from <https://phys.org/news/2023-04-recycling-plastic-packaging-fragrances.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--