

Timing of snowshoe hare winter color swap may leave them exposed in changing climate, study finds

April 12 2023, by Samantha Murray



A mismatched snowshoe hare in winter coat in a snow-free landscape in Kluane, Yukon, Canada. Pure white winter coat against snow-free landscape makes snowshoe hares easily detectable by predators, which can reduce their survival during the winter. The study found, however, that hares that wait longer to turn white in autumn face the greatest chance of being eaten by predators during the winter. Credit: Lance Goodwin



Like many animals in the far north, snowshoe hares change their coats from brown to white each autumn. Come winter, these all-white hares are harder for predators to spot against the snowy landscape, helping ensure their survival.

But as <u>global warming</u> reduces snowfall in the region, will the hares' seasonal color swap continue to keep them safe?

Scientists from the University of Florida and Canada are beginning to answer that question. Published in *Proceedings of the Royal Society B*, Their new study, which used 44 years of data on snowshoe hares in Canada's Yukon Territory, shows that as the region has warmed, hares are waiting longer to turn pure white.

However, the data also suggest that waiting too long can be deadly. The scientists found that hares that were more brown than white in autumn were less likely to survive the winter

"When it comes to the switch from brown to white, which we call molting, timing is everything. We can see in this study that <u>climate</u> <u>change</u> is making it harder for the hares to get that timing just right." said Madan Oli, first author of the study and a professor in the UF/IFAS wildlife ecology and conservation department.

"Hares begin to change color based on environmental cues such as temperature and snow depth, but the change doesn't happen overnight. The whole molting process takes about a month. So, if hares are beginning to molt later in the year, there is a greater chance they will not be fully white by the time their environment is white. These hares are then more vulnerable to predators."





Snowshoe hare in winter coat in a snowy landscape in Kluane, Yukon, Canada. Pure white winter coat provides camouflage against snow, which reduces risk of being seen and caught by predators. Credit: Alice Kenney

Snowshoe hares are what ecologists call a keystone species because so many other animals depend on them for food, said Charles Krebs, senior author of the study and a professor emeritus at the University of British Columbia.

If the snowshoe <u>hare</u> population takes a nosedive, this could affect the entire ecosystem, he said.

"Practically everything eats snowshoe hares in the <u>boreal forest</u>. A big change in the number that survive the winter impacts their ability to replenish themselves over the summer breeding season, which could



disrupt the whole food web," Krebs said. The boreal forest, or taiga, rings the norther part of the globe and includes parts of Alaska, Canada, Europe and Asia.

Snowshoe hares are among 21 bird and <u>mammal species</u> in boreal forest that turn white for winter, which scientists call seasonal molting. This study is the first to track how seasonal molting has shifted with rising <u>global temperatures</u>.

The decades of data collection by Krebs and colleagues also makes the study unique, Krebs said.

"When you look over a forty-four-year span, you can really document a pattern. This study demonstrates the value and importance of long-term ecological monitoring," Krebs said.



Brown snowshoe hare. Hares turn brown for the summer when the snows recede.



Credit: Alice Kenney

Between 1977 and 2021, Krebs and his collaborators went to the Kluane National Park and Reserve in Canada's Yukon Territory each autumn and spring to study snowshoe hares. In the remote, often extremely cold wilderness they trapped snowshoe hares, recorded the percentage of white and brown in each hare's coat, tagged each hare (or noted if it was already tagged) and set it free.

The researchers recorded observations for nearly 4,500 hares, with some caught multiple times over the years. Oli, who specializes in evaluating <u>animal populations</u> from a statistical and modeling perspective, and then wildlife ecology doctoral candidate Vratika Chaudhary, analyzed this massive dataset. The current study presents the results of that analysis.

The team isn't done studying snowshoe hares in the Yukon, Krebs said. The scientists are currently recording the hare coat color and behavior using motion activated cameras. Images captures by these cameras will show how much—or how little—the hares coats are helping them match their environment in real time.

More information: Madan K. Oli et al, Does coat colour influence survival? A test in a cyclic population of snowshoe hares, *Proceedings of the Royal Society B: Biological Sciences* (2023). DOI: 10.1098/rspb.2022.1421

Provided by University of Florida

Citation: Timing of snowshoe hare winter color swap may leave them exposed in changing



climate, study finds (2023, April 12) retrieved 28 June 2024 from <u>https://phys.org/news/2023-04-snowshoe-hare-winter-swap-exposed.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.