


Astronomers claim to have solved mystery of the runaway supermassive black hole

May 9 2023



Above: Image of the object observed with the Hubble Space Telescope. It shows the emission in the ultraviolet part of the spectrum. Middle: Ultraviolet image of a local galaxy without a bulge and observed edge-on (IC 5249). The similarities are obvious. Bottom: The same galaxy IC 5249 observed in the visible part of the spectrum. The spatial scales of the three images are identical. Credit: ESA/Hubble

A mysterious trail of stars formed eight thousand million years ago and recently discovered by the Hubble Space Telescope has been a challenge to a number of research groups. Its size is similar to that of the Milky Way, and this very long narrow structure has given rise to several explanations of its origin.

According to a controversial initial hypothesis, this trail of stars could be the result of the passage of a supermassive black hole through a huge cloud of gas. This idea quickly fired the imagination of the astronomical community, because it needs a large set of complex exceptional circumstances. For this reason several scientific teams have continued to explore different and less exotic scenarios that could explain the observations.

In a recent study published in *Astronomy & Astrophysics*, researchers at the Instituto de Astrofísica de Canarias (IAC) have reached the conclusion that this unusual structure of stars could be interpreted as a galaxy without a bulge seen edge-on. Galaxies of this type, also called thin or flat [galaxies](#), are relatively common.

"The motions, the size, and the quantity of stars fits what has been seen in galaxies within the local universe," explains Jorge Sanchez Almeida, an IAC researcher who is the first author of the article. "It's a relief to have found the solution to this mystery; the new proposed scenario is much simpler. In one sense it is also a pity, because the existence of fleeing [black holes](#) is expected, and this could have been the first one to be observed."

To support the hypothesis of the interpretation in terms of a galaxy, the team compared the mysterious structure with a well-known local galaxy without a bulge, IC5249, which has a similar mass of stars, and found surprising agreement. In the words of Mireia Montes, an IAC researcher who is a co-author of the article, "When we analyzed the velocities of

this distant structure of stars, we realized that they were very similar to those obtained from the rotation of galaxies, so we decided to compare a much closer galaxy, and found that they are extraordinarily similar."

"We also looked at the relation between the mass of the assumed galaxy and its maximum velocity of rotation, and discovered that indeed it is a galaxy which behaves like a galaxy," states Ignacio Trujillo, an IAC researcher who took part in the study. "It is an interesting object, because it is quite a large galaxy at a very large distance from Earth, where the majority of the galaxies are smaller," he adds.

Upcoming observations will allow the study of this object in greater detail.

More information: Jorge Sánchez Almeida et al, Supermassive black hole wake or bulgeless edge-on galaxy?, *Astronomy & Astrophysics* (2023). [DOI: 10.1051/0004-6361/202346430](https://doi.org/10.1051/0004-6361/202346430)

Provided by Instituto de Astrofísica de Canarias

Citation: Astronomers claim to have solved mystery of the runaway supermassive black hole (2023, May 9) retrieved 5 December 2023 from <https://phys.org/news/2023-05-astronomers-mystery-runaway-supermassive-black.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.