

Study establishes molecular basis for interaction between essential protein complex and its regulator

May 25 2023, by Leah Mann



Credit: CC0 Public Domain

The labs of Lauren Jackson, associate professor of biological sciences and biochemistry, and Todd Graham, Stevenson Professor of Biological Sciences at the College of Arts and Science and professor of cell and developmental biology, recently published a study in the *Journal of Cell Biology* describing a significant interaction between an essential protein complex used for protein and lipid transport—the COPI complex—and



its regulator protein.

The authors include Betty Xie and Cameron Cohen, two graduate students in the Jackson lab, and Swapneeta Date, senior scientist in the Graham lab. Together, the two groups established how COPI interacts with the regulator <u>protein</u> Glo3, a member of the ArfGAP protein family of enzymes. Disruptions of the interaction between COPI and Glo3 lead to deleterious effects in cells.

We sat down with Jackson to find out more about this research.

What issue/problem does your research address?

Our lab studies the cellular "FedEx system" that allows proteins and lipids to move between membrane-bound compartments. This work focused on COPI, which is required in all eukaryotes. Mutations in COPI are associated with multiple human diseases, including cancers and microcephalies, which are conditions in which a patient's head is abnormally small. Pathogens, including viruses like SARS-CoV-2, also hijack COPI to take over cells.

While the field has previously established that molecules in the ArfGAP family affect COPI function, the mechanisms involved in this interaction have not been well understood. We sought to determine the basis for the interaction between COPI subunits and a specific ArfGAP protein, Glo3.

What was unique about your approach to the research?

We undertook a multi-disciplinary approach to address these questions using biochemistry, biophysics, computational artificial intelligence-



based modeling, and in vivo approaches including fluorescence imaging in budding yeast. This combination of approaches allows us to use the molecular information generated from biochemical, biophysical, and computational studies to design hypotheses that could be used to test function directly in budding yeast, our model organism.

What were your findings?

We discovered a molecular "code" that explains how COPI recruits ArfGAP proteins. We identified specific amino acid residues on each protein that mediate the interaction between COPI and ArfGAP. We then tested how breaking the interaction affected the movement of transmembrane protein cargo using budding yeast as our model organism. Breaking this interaction specifically disrupted the shape of a key organelle called the Golgi, and it also misdirected specific cargo proteins to the wrong cellular location.

What do you hope will be achieved with the research results in the short and long terms?

Overall, many of these proteins are strongly conserved across species, which suggests our results will be relevant to human biology. This study provides a variety of molecular tools that we can use to dissect these important cellular events that drive essential membrane trafficking—the process of moving proteins and other molecules throughout the cell using membrane-bound vesicles.

What are the benefits of this research?

This research contributes to our understanding of fundamental biological processes in cellular sorting. Obtaining <u>molecular information</u> allows us to probe complex cellular sorting pathways so that we can understand



human physiology and what goes wrong in disease states, such as in neurological disorders or cancers, when essential proteins are mutated or lost.

Where is this research taking you next?

This work builds on our ongoing collaboration with Todd Graham's lab. Together, the Jackson and Graham labs use a variety of methods to understand how the COPI/Glo3 interaction drives specific cellular sorting events (Jackson lab) and how the interaction is affected by posttranslational modifications and sorting of SNAREs, a family of proteins that play essential roles in vesicle fusion events.

More information: Boyang Xie et al, An interaction between β' -COP and the ArfGAP, Glo3, maintains post-Golgi cargo recycling, *Journal of Cell Biology* (2023). DOI: 10.1083/jcb.202008061

Provided by Vanderbilt University

Citation: Study establishes molecular basis for interaction between essential protein complex and its regulator (2023, May 25) retrieved 29 June 2024 from https://phys.org/news/2023-05-molecular-basis-interaction-essential-protein.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.